

Agenda Item 7

Review of the Conservation Plan

Document 7

**ASCOBANS Conservation Management
Plan (CMP) for Harbour Porpoise
(*Phocoena phocoena*) in the
Greater North Sea – Draft**

Action Requested

Provide feedback

Submitted by

Secretariat



ASCOBANS
Conservation Management Plan
(CMP)

For Harbour Porpoise
(*Phocoena phocoena*)
in the Greater North Sea



February 2023

Table of Contents

Executive Summary	4
Summary of actions	5
1. Introduction	6
1.1. Necessity for a Conservation Management Plan	6
1.2 Overall objective of the Conservation Management Plan	6
1.3. Development of the Conservation Management Plan.....	7
1.4. Coordination and Governance of the Conservation Management Plan	7
2. Legal framework	8
3. Biology and status of harbour porpoise	8
3.1. Summary of biology and ecology	8
3.2. Abundance and distribution	13
4. Pressures	19
4.1. Summary of pressures	19
4.2. Attributes of the population for monitoring, mitigation and research	23
5. Conservation Status	24
5.1. Critical Habitats	25
5.2. Dealing with inadequate data.....	27
6. Actions	28
6.1. Summary of actions	28
6.2. Actions and Tasks	29
Action RES-01: Identify the priority issues and relevant stakeholders	30
Action RES-02: Improve estimates of bycatch rates to support development of conservation strategy	31
Action MIT-01: Implement and assess pinger and other mitigation measures to reduce bycatch	32
Action MON-01: Implement a wide-scale surveillance programme to monitor trends in distribution and abundance in the Greater North Sea	33
Action RES-03: Improve understanding of causes of seasonal and annual variation in abundance and distribution, particularly in relation to human activities and environmental change, to facilitate the consideration of the species within marine spatial plans	34
Action MON-02: Monitoring of health and nutritional status, diet, life history parameters and causes of mortality.	35
Action RES-04 Further our understanding of population structure	36
Action MIT-02: Improve understanding of and develop mitigation for the risks of anthropogenic sound	37
Action MON-03: Ensure screening and assessment of the occurrence and effects of hazardous substances.....	38
Action MON-04: Monitor for potential increases in anthropogenic activities that lead to	

incidences of death, injury or adverse health effects including cumulative effects.....	39
Action MON-05: Monitor habitat quality, including protected sites, to ensure management is effective and that the ecological functions are maintained.....	40
7. Public awareness and capacity building	41
7.1 Public awareness tasks	41
Annex 1: International conventions and agreements	42
Annex 2: UK and Norway National Legislation	48

Summary of actions

Priority	Action	Code
Essential	Identify the priority bycatch issues and relevant stakeholders	RES-01
Essential	Improve estimates of bycatch rates to support development of conservation strategy	RES-02
Essential	Implement and assess pinger and other mitigation measures to reduce bycatch	MIT-01
High	Implement a wide-scale surveillance programme to monitor trends in distribution and abundance in the Greater North Sea	MON-01
High	Improve understanding of causes of seasonal and annual variation in abundance and distribution, particularly in relation to human activities and environmental change, to facilitate the consideration of the species within marine spatial plans	RES-03
High	Monitor health and nutritional status, diet, life history parameters, and causes of mortality	MON-02
Medium	Further our understanding on population structure	RES-04
Medium	Improve understanding of and develop mitigation for the risks of anthropogenic sound	MIT-02
Medium	Ensure screening and assessment of the occurrence and effects of hazardous substances	MON-03
Low	Monitor for potential increases in anthropogenic activities that lead to incidences of death, injury or adverse health effects	MON-04
Low	Monitor habitat quality, including protected sites, to ensure management is effective and that the ecological functions are maintained.	MON-05

ASCOBANS Conservation Management Plan (CMP) for Harbour Porpoise (*Phocoena phocoena*) in the Greater North Sea

1. Introduction

1.1. Necessity for a Conservation Management Plan

Harbour porpoises (*Phocoena phocoena*, Linnaeus 1758) are widely distributed in shelf waters of the temperate North Atlantic and of the North Pacific Oceans and in some semi-enclosed seas, such as the Black and Baltic Seas. The Greater North Sea is an important habitat for harbour porpoises in the North East Atlantic, a region where a large percentage of the North-east Atlantic population inhabits. Harbour porpoises are exposed to a number of anthropogenic pressures, of which bycatch in commercial fisheries is considered the greatest threat. The mobile nature of both the species and the key threat means a conservation management plan would provide the most effective way to achieve the conservation objectives.

International collaboration on the conservation and management actions developed in this plan will be necessary. OSPAR has listed the harbour porpoise as a threatened and declining species, with a focus on tackling bycatch. Support by both ASCOBANS and OSPAR will be key, and will require co-operation by many stakeholders, ranging from local and national governments, through intergovernmental bodies to industry and NGOs. This CMP follows the International Whaling Commission (IWC) conservation management plan template, also adopted by ASCOBANS. This should be considered a dynamic document and that changes will be undertaken periodically through an expert review process to enable the development of new or modified actions as appropriate.

1.2 Overall objective of the Conservation Management Plan

A conservation management plan must have measurable objectives by which its success or failure can be evaluated regularly, and to ensure that required changes are identified and actioned promptly. Failure to monitor progress will result in inaction and subsequent failure of the CMP. Integral and essential to the plan are, therefore, monitoring of:

- Regional and overall trends in the North Sea harbour porpoise management unit;
- human activities identified to pose potential risk to the species;
- implementation of mitigation measures and;
- the assessment of effectiveness of those measures.

ASCOBANS intermediate conservation objective aims to '*restore and/or maintain biological or management stocks of small cetaceans at the level they would reach when there is the lowest possible anthropogenic influence*' with '*a suitable short-term practical sub-objective to restore and/or maintain stocks/populations to 80% or more of the carrying capacity*' (ASCOBANS, 1997). To work towards achieving this intermediate goal, the CMP identifies the key pressures and threats facing the management unit, gaps in evidence and information, and proposes actions necessary to achieve the goal of maintaining the management unit and population at a favourable conservation status. These actions include coordination of monitoring programmes on direct and indirect pressures, including bycatch, marine pollution and anthropogenic noise, to allow a full assessment of the effects on the management unit. The

actions in this CMP also complement and support wider measures for small cetaceans in the North East Atlantic.

1.3. Development of the Conservation Management Plan

The North Sea harbour porpoise Conservation Management Plan will be coordinated under a hierarchical structure clearly outlining roles and responsibilities (Figure 1), designed to ensure effective implementation. A Steering Group (SG) has been formed to drive implementation of the plan. Co-operation and complementarity with the work of other ASCOBANS working groups will be sought. Of particular relevance are the ASCOBANS/ACCOBAMS Joint Bycatch Working Group, ASCOBANS Pollution Working Group, ASCOBANS Working Group on Resource Depletion, the Joint ACCOBAMS/ASCOBANS Working Group on the Marine Strategy Framework and the ACCOBAMS/ASCOBANS Joint Noise Working Group.




Figure 1: CMP communication structure.

1.4. Coordination and Governance of the Conservation Management Plan

The coordinator and SG will ensure cooperation between all stakeholders including national governments in the NE Atlantic, European Commission, intergovernmental organisations including fisheries management authorities, ICES and OSPAR, Advisory Councils and other relevant bodies, such as NGOs, universities and institutes, and appropriate industry representatives. Their role specifically is to encourage countries to harmonise their national efforts, including allocation of funding. The CMP will be a dynamic document and subject to regular revision to ensure the information remains current.

To ensure efficiency and to drive the plan forward, the following tasks have been identified:

1. The SG has appointed a coordinator (or chair) to oversee implementation of the plan. The SG and coordinator will together:
 - develop and maintain the Terms of Reference for the SG to ensure that the actions are implemented;
 - coordinate and drive the implementation of the CMP (including assessing funding options where appropriate) and promote the CMP to relevant stakeholders;
 - collate reports on the progress of implementation, effectiveness and issues encountered and report annually to the Advisory Committee on the progress of the CMP, establish further implementation priorities and make appropriate recommendations;

-
- encourage cooperation between ASCOBANS and Range States.
2. ASCOBANS Parties and Range States will report annually on implementation of the CMP.
 3. The coordinator/SG will evaluate the CMP every six years and amend the document where required as agreed by the Advisory Committee.

2. Legal framework

The 5th International Conference for the Protection of the North Sea (Bergen, Norway, 20-21 March 2002) called for a recovery plan for harbour porpoises in the North Sea to be developed and adopted (Paragraph 30, Bergen Declaration). This was subsequently taken forward by ASCOBANS with the publication of the Conservation Plan for Harbour Porpoises (*Phocoena phocoena* L.) in the North Sea (ASCOBANS 2009).

There is a broad list of drivers behind harbour porpoise conservation which aim to address all aspects of anthropogenic impact on the species, either specifically for harbour porpoise, or as part of a wider strategy for cetaceans or marine mammals. A summary of the legal framework relevant to harbour porpoise including conventions and agreements can be found in Annex 1. On 31 January 2021, the UK left the European Union whilst remaining a Party to ASCOBANS and OSPAR. Annex 2 contains the relevant national legislation for the UK pertaining to cetacean conservation. As a range state to the ASCOBANS area, a summary of Norway's national legislation has also been included in Annex 2.


3. Biology and status of harbour porpoise

3.1. Summary of biology and ecology

Within North eastern Atlantic waters, harbour porpoise comprises a continuous biological population that extends from the Northwest African coast northwards to the Arctic waters of Norway and Iceland (Fontaine et al. 2007, Fontaine et al. 2010, Fontaine et al. 2014, Evans 2020). Separate (sub-) populations have been recognised in the Belt Sea and Baltic Sea proper (Wiemann et al. 2010, Galatius et al. 2012, Sveegaard et al. 2015, Lah et al. 2016), whereas, a separate sub-species *Phocoena phocoena meridionalis*, of a larger-sized morphotype (Donovan and Bjorge 1995) (Donovan and Bjorge, 1995), has been proposed to occur in Iberian and Mauritanian waters (Jung et al. 2009, Fontaine et al. 2014, Fontaine et al. 2017, Murphy et al. 2020)

A comprehensive review of the status of harbour porpoises in the entire Atlantic Ocean was conducted in 2018 (IMR-NAMMCO 2019) which considered a range of parameters for defining assessment areas including population structure, as well as measurements of time-integrated ecological tracers and morphological differences, ICES areas/divisions boundaries, and the spatial extent of human activities (ICES WGMME 2013, 2014, IAMMWG 2015, OSPAR 2017, Murphy et al. 2020, Chehida et al. 2021). OSPAR and Contracting Parties have adopted the proposed IMR-NAMMCO assessment areas as assessment units for reporting under the Marine Strategy Framework Directive (MSFD) (Figure 2). For the current CMP, these boundaries will be employed for the Greater North Sea harbour porpoise management unit, and is defined as ICES divisions IVa, b, c, VIId and the northern part of IIIa.

Figure 2: Map of the assessment areas for harbour porpoise in the North Atlantic proposed by IMR/NAMMCO and adopted by OSPAR, with the ICES fishing areas super-imposed. (From IMR/NAMMCO, 2019. Abundance estimates for the assessment units are available in the online supplementary files: <https://nammco.no/topics/scientific-workshops-symposia-reports/#2018>).



The harbour porpoise is one of the smallest toothed whales (typically less than 1.6m in size in the Greater North Sea) that inhabits coastal and continental shelf waters, including those of the ASCOBANS agreement area (Evans 2020). The lifespan of harbour porpoises varies between populations and geographic areas, although not significantly between sexes. They live for notably less time than most other marine mammal species, with an average lifespan of 8-13 years (Lockyer 2003). This shorter lifespan increases the sensitivity of the harbour porpoise population growth rate to fluctuations in other factors such as juvenile mortality or reproductive rates.

The Joint IMR/NAMMCO International Workshop on the Status of Harbour Porpoises in the North Atlantic undertook a full status review of harbour porpoises in the Greater North Sea in 2019 (Murphy et al. 2019). A summary of such is presented herein, along with more recent work, focusing on elements of biology and ecology. While a number of historical and contemporary life history studies have been undertaken within the region, assessments at the scale of the management unit is lacking. A basin wide assessment would substantially increase the statistical power of such studies, through increasing sample sizes of all age-sex groups.

Murphy et al. (2020) assessed demographic characteristics and determined key biological parameters in stranded and bycaught male and female harbour porpoises within the UK North Sea, including an assessment of temporal variation in those parameters (Table 1). Harbour porpoises in the Greater North Sea management unit are significantly smaller in body size, both at attainment of sexual and physical maturity compared to porpoises within the Celtic and Irish Seas management unit (which incorporates waters off western France; see Figure 2)

(Murphy et al. 2020). The Celtic and Irish Seas management unit is viewed as a mixing or transition zone between the North-east Atlantic population and the larger morphotype Iberian sub-species (Murphy et al. 2019).

Table 1. Asymptotic length and age estimated using the Gompertz growth model, Length at 50% maturity (L50), and Age at 50% maturity (A50) for female and male harbor porpoises in the North Sea Management Unit (MU) and Celtic and Irish Seas MU sampled in UK waters for two time periods, 1990–1999 and 2000–2012. Adapted from Murphy et al. (2020).

MU	Sex	Time period	Asymptotic length (cm)	Asymptotic age (yrs)	L50 (cm)	A50 (yrs)
North Sea	Females	1990–1999	155.37	7.21	138.90	3.8
		2000–2012		11.66	139.18	4.8
	Males	1990–1999	140.94	5.72	133.27	3.56
		2000–2012		7.62	129.47	3.62
Celtic & Irish Seas	Females	1990–1999	162.94	7.21	146.56	3.8
		2000–2012		11.66	146.94	4.8
	Males	1990–1999	146.50	5.72	138.73	3.56
		2000–2012		7.62	133.46	3.63

Female and male harbour porpoises in the UK North Sea were reported to attain physical maturity at 155.4 cm, and 140.9 cm, respectively (Murphy et al. 2020). An increase in the age at asymptotic length was observed in both sexes over the last few decades, along with a significant decline in the Gompertz growth rate parameter that was more apparent in the female data (Murphy et al. 2020). Females also significantly increased in their average age at attainment of sexual maturity (A50), while this parameter remained relatively stable for males (Table 1). Male harbour porpoises however, significantly declined in the average length attained at sexual maturity (L50), while no significant difference was observed for females. It was suggested that availability of suitable prey resources could possibly be a limiting factor and an explanation for the observed results, though a combination of other factors was not ruled out (Murphy et al. 2020). For harbour porpoises inhabiting the German North Sea, Kesselring et al. (2017) also reported that females attained sexual maturity at an older age (of 4.95 years) and were also dying, on average, at an older age than Baltic porpoises (5.70 (\pm 0.27) years vs 3.67 (\pm 0.30) years, respectively). It is estimated that only 54.66% of females in German North Sea would participate in reproduction (Kesselring et al. 2017).

Looking at contemporary reproductive rates for the Greater North Sea, Murphy et al. (2020) reported a reduced reproductive rate (29% pregnancy rate) in porpoises inhabiting UK waters, though the sample was heavily biased towards stranded animals that died of infectious disease, or other causes such as starvation. Earlier work using all available UK data estimated a pregnancy rate of 50% for a control group of 'healthy' females – females that died of traumatic causes of death such as by-catch, boat/ship strike, bottlenose dolphin attacks or dystocia - a pregnancy rate almost half that reported for other geographical regions, and a notable increased incidence of reproductive pathologies, including reproductive failure, potentially associated with exposure to endocrine disrupting chemicals (Murphy et al. 2015). While a recent study on harbour porpoises in Dutch waters using samples obtained between 2006 and 2019 reported that maternal nutritional status had a significant effect on foetal size, and females in poor health had a lower probability of being pregnant and were less likely to carry a foetus to term (IJsseldijk et al. 2021). Within the

study, a pregnancy rate of 28% (51 of 180 mature females) was determined for the whole sample and a higher pregnancy rate of 58% (22 of 38 mature females) was determined for 'healthy' females, while an average age at sexual maturity was estimated at 4 years for females (Ijsseldijk et al. 2021b).

Harbour porpoises are opportunistic piscivore predators, with diet varying significantly according to prey availability (Murphy et al. 2019, Lambert 2020). A huge variety of prey taxa have been recorded from the stomachs of stranded harbour porpoise in the Greater North Sea. However, the diet of individuals tends to be dominated by 2-4 species at any one time (Pierce et al. 2007). The most commonly identified key prey groups are gadoids (mostly whiting), gobies, sandeels and clupeids (both herring and sprat) (Lambert 2020, Pierce et al. 2022). Table 2 includes a summary of studies assessing the diet of porpoises in the Greater North Sea.

Though the species is noted to be a generalist, it may be vulnerable to the depletion of key prey species, impacting survival and reproduction (MacLeod et al. 2007, Leopold 2015, Pierce et al. 2022), as their high metabolic rate (large surface body to volume ratio) requires efficient foraging including ultra-high capture rates of high energy density prey (Wisniewska et al. 2016, Wisniewska et al. 2018). Changing prey dynamics may have been responsible for the re-distribution of porpoises within the Greater North Sea since the 1990s, with the decline in sandeels (*Ammodytes marinus*) in the northern North Sea, and the re-invasion of the southern North Sea by sardine (*Sardina pilchardus*) (Mahfouz et al. 2017). Though, consumption of leaner gadoids and gobies were also reported in the stomachs of porpoises inhabiting the southern North Sea, animals that were in poorer body condition, associated with emaciation/starvation (Leopold 2015). Potential impacts from resource depletion is discussed in Annex 3.

Table 2. Harbour porpoise diet inferred from stomach content analysis in the southern North Sea and adjacent areas (n = number of stomachs).

Area (year of stranding)	n	Main prey	Reference
Southern North Sea (2010–2013)	14	Gobies, whiting, sandeel	(Mahfouz et al. 2017)
Germany	34	Sandeels, sole	(Benke and Siebert 1996)
Germany	36	Sole, cod	(Lick 1991)
Belgian coast (1997–2011)	64	Gobies, sandeels, whiting, <i>Trisopterus</i> sp.	(Haelters et al. 2012)
Belguim coast (1997-2018)	180	Whiting, gobies, sandeels , herring and sprat	(Lambert 2020)
Dutch coast (2006)	64	Gobies, sandeels, sprat, herring, whiting, twait, shad	(Leopold and Camphuysen 2006)
Dutch coast (2003-2010)	229	Whiting, gobies, sandeels, sprat, herring, cod	(Leopold et al. 2011)
Dutch coast (2003-2010)	76	Whiting, gobies, lesser sandeels, sprat, herring, cod	(Jansen 2013)Jansen (2013)
Dutch coast (2003-2014)	600	Adults: gadoids, clupeids and sandeels Juveniles: Gobies	(Schelling et al. 2014)
Dutch coast (2006-2014)	824	Gadids, gobies, sandeels, clupeids	(Leopold 2015)
English Channel (1998–2003)	7	Pouting, gobies	(De Pierrepont et al. 2005)
East Scotland (1959–1971)	93	Herring, sprat, whiting	(Rae 1965, Rae 1973)
Scotland (1992-1996)	72	Whiting, sandeels, herring	(Santos 1998)
Scotland (1992–2003)	188	Whiting, sandeels, gadids, <i>Trisopterus</i> sp.	(Santos et al. 2004)
UK (1989–1994)	100	Gadids, sandeels, gobies	(Martin 1996)
Denmark, Sweden, Norway	197	Herring, gadids	(Aarefjord et al. 1995)

Analysis of stranding records for the North Sea between 1990 and 2017 revealed that incidences of strandings have increased, with a sharp rise observed within the southern North Sea since 2005, and a higher density of neonatal strandings in the eastern North Sea (Ijsseldijk et al. 2020)(see Figure 3). Incidences of unusual mortality events have also been reported in Danish (Wright et al. 2013), and Dutch waters (Ijsseldijk et al. 2020, IJsseldijk et al. 2021a). Harbour porpoises in the German North Sea have been assessed to be in a poor general health status, with a higher incidence of severe pathological lesions, especially of the respiratory tract (with pneumonia considered the most common cause of death), compared to harbour porpoises inhabiting more northern waters, which may be due to exposure to chemical pollutants (Siebert et al. 2006, Siebert et al. 2009) (see Annex 3). While infectious disease may be the leading cause of death, fisheries interactions was the leading direct anthropogenic cause of death in porpoises stranded along Dutch and UK coastlines, with a lower number of cases of trauma due to vessel strikes being reported (Deaville and Jepson 2011, Deaville 2016, 2018, IJsseldijk et al. 2022). Though incidences of trauma resulting from inter-species interactions have been reported at a higher occurrence, notably from bottlenose dolphins (Deaville and Jepson 2011, Deaville 2018), as well as grey seals (Leopold et al. 2015, IJsseldijk et al. 2022). In the light of recent work demonstrating acoustic trauma in porpoises due to explosions in the Baltic Sea (Siebert et al. 2022), the North Sea group has expressed concern over similar activities occurring in the North Sea, as surviving animals might have impaired hearing which, among other things, could affect their ability to detect nets and find prey. While, cases of starvation/emaciation have been increasing in recent years among necropsied porpoises in (all) UK waters (Deaville 2018), and was also a leading cause of death among porpoises that died during an unusual mortality event in Dutch waters in 2011 (Ijsseldijk et al. 2022).




Figure 3. Study area showing the density of all recorded harbour porpoise strandings over three time periods (Taken from (Ijsseldijk et al. 2020)).

3.2. Abundance and distribution

Abundance and occurrence of harbour porpoises have fluctuated over the last 100 years within the North-east Atlantic. A decline in both strandings and observations occurred in the southern North Sea, English Channel and French Atlantic coasts from the 1950s onwards (Smeenk 1987, Evans 1992, Addink and Smeenk 1999, Camphuysen 2004, Evans et al. 2008, Jung et al. 2009). Within the last two decades, porpoises started to return again to these waters, which included a re-distribution of animals from the northern to the southern North Sea, as well as the re-population of central English Channel and waters off the French Atlantic coast (Camphuysen 2004, Hammond et al. 2013, Hammond et al. 2017, Laran et al. 2017).

Abundance estimates and trends are a key parameter in any population assessment and reliable estimates are required for sound scientific management of stocks. Methods have been developed for surveys targeting small cetaceans (e.g. SCANS) that have resulted in robust estimates of harbour porpoise abundance (Hammond et al. 2013). To date, four SCANS surveys have been undertaken that include the North Sea (1995, 2005, 2016 and 2022; Figure 4). From these it has been estimated that the population of harbour porpoise in the North Sea has been stable (potentially increasing very slowly) since 2005 (Figure 5).

The distribution of harbour porpoises is not static in space or time. For instance, in records from 1979-1997, sighting rates in the south eastern North Sea, the southern Bight and the northern English Channel were substantially lower than in areas further north (Evans et al. 2003, Reid et al. 2003). Thereafter, surveys reported higher sighting (Brasseur et al. 2004, Scheidat et al. 2004) and strandings rates (Haelters et al. 2002, Jauniaux et al. 2002, Camphuysen 2004, Kiszka et al. 2004) in the southern North Sea and southern Bight. This increase in both sighting and stranding rates in these southern parts of the North Sea over a relatively short period of time suggested a redistribution of animals from other areas rather than a sudden and rapid increase in population growth in the southern North Sea. This redistribution appears to have been maintained in subsequent SCANS surveys (2013 and 2022).

The most robust modelling of the distribution of harbour porpoise in the North Sea is by Gilles et al. (2016), who generated modelled distributions for the period 2005-2013 for spring, summer and autumn, a period after the main re-distribution of the species within the Greater North Sea (Figures 6-8). The predicted distributions for all three seasons show higher density in the western North Sea off the coast of the UK and lower densities in the eastern North Sea closer to Denmark and Germany. In summer, the predicted higher density area appears to extend slightly further south in summer than in autumn and spring (Gilles et al. 2016).




Figure 4: Density surface maps from the SCANS surveys 1994 (top left), 2005 (top right), and 2016 (bottom left), and sightings from 2022 (bottom right) (Adapted from (Hammond et al. 2013, Lacey et al. 2022) and SCANS IV, unpublished).




Figure 5: Trend in harbour porpoise abundance for the North Sea in 1994, 2005 and 2016. Estimated rate of annual change is 0.8% (95% confidence interval: -6.8% to 9.0%), P = 0.18 (Adapted from Hammond et al., 2017).




Figure 6: Predicted harbour porpoise densities in the North Sea in spring (March-May) 2005-2013. Upper panel: The overlaid contours are associated jackknife standard deviations (SD). The black and white dashed boundary depicts the sampling coverage in spring. Lower panel: Lower and upper lognormal 90% confidence intervals of predicted density. From Gilles et al. (2016).




Figure 7: Predicted harbour porpoise densities in the North Sea in summer (June-August) 2005-2013. Upper panel: The overlaid contours are associated jackknife standard deviations (SD). Lower panel: Lower and upper lognormal 90% confidence intervals of predicted density. From Gilles et al. (2016).




Figure 8: Predicted harbour porpoise densities in the North Sea in autumn (September-November) 2005-2013. Upper panel: The overlaid contours are associated jackknife standard deviations (SD). The black and white dashed boundary depicts the sampling coverage in spring. Lower panel: Lower and upper lognormal 90% confidence intervals of predicted density. From Gilles et al. (2016).

4. Pressures

4.1. Summary of pressures

As harbour porpoise occurs throughout the European continental shelf waters the species can be affected by a range of human activities occurring in the same waters (IAMMWG 2015, ICES WGMME 2015, IMR-NAMMCO 2019, WGMME 2019). A detailed summary of information on pressures including evidence gaps, can be found in Annex 3.

The single most significant anthropogenic threat to harbors porpoise is bycatch in bottom-set static nets (Read et al. 2006, Bjørge et al. 2013, Scheidat et al. 2013, ICES Advice 2014, Nabe-Nielsen et al. 2014, van Beest et al. 2017, FAO 2018, Northridge 2018, STECF 2019, Evans et al. 2021, ICES Advice 2021b, Moan and Bjørge 2023). The ICES Workshop on estimation of MOrtality of Marine MAMmals due to Bycatch (WKMOMA) in 2021 addressed the special request from OSPAR regarding the bycatch mortality in marine mammals, including harbour porposies in the Greater North Sea (ICES 2021). The workshop was tasked with generating bycatch rates (e.g. specimens per day at sea) and associated confidence intervals for static and towed gears (at least Métier Level 4) (Table 3), in addition to generating assessment unit and métier specific bycatch mortality estimates (Table 4). Highest bycatch rates were observed in gillnet metiers, particularly those deployed from large vessels over the period 2015 to 2020. Effort data from Norwegian vessels and small German vessels were not available for the assessment. Further, there was a potential bias in the dataset as for one member country, vessels with high bycatch rates were targeted for monitoring, increasing bycatch rates by a factor of up to 5 in set gill nets (GNS) and drift nets (GND), and 3.5 in trammel nets (GTR) (ICES Advice 2021b).

Other anthropogenic activities that may affect harbour porpoise include:

- underwater noise (e.g. as generated by shipping, pile driving, seismic surveys, detonation of explosives and acoustic deterrent devices) (Todd et al. 1999, Stone and Tasker 2006, Bailey et al. 2010, Brandt et al. 2011, Dähne et al. 2013, Dähne et al. 2017, Stone et al. 2017, Wisniewska et al. 2018, Roberts et al. 2019));
- pollution (particularly persistent organic pollutants such as polychlorinated biphenyls [PCBs]) (Jepson et al. 2005, Murphy et al. 2015, Jepson et al. 2016, Williams et al. 2020, Williams et al. 2023),
- collision risks (IAMMWG 2015, Robbins 2022)
- prey depletion (Santos and Pierce 2003, Pierce et al. 2022),
- marine debris (Unger et al. 2017); and
- environmental change (Learmonth et al. 2006, IAMMWG 2015).

A summary of pressures, related activities, and current levels of evidence for pressures associated with harbour porpoise is presented in Table 5. These are based on the threat matrices (ICES WGMME 2015, WGMME 2019). The pressures have been split into the following categories after Authier et al. (2017):

- **Primary** (direct mortality);
- **Secondary** (health degradation, with indirect effect on demography) and;
- **Tertiary** (behavioural disruption, with indirect effect on health and therefore demography).

Some pressures are identified as medium or low priority in terms of action required when assessed in isolation. However, it should be noted that when acting in combination with other pressures, the risk to the species could increase. A strategic approach to conservation should be considered to account for the cumulative impacts of non-lethal (secondary and tertiary) pressures acting on the individuals and the combined demographic effects of all pressures on the population.

Table 3. The bycatch rate per ICES subarea and métier level 4 for the harbour porpoise Greater North Sea AUs with data from 2015 to 2020. Both the estimated frequency of bycatch events and the estimated number of individuals per bycatch event is shown. (GNS = gill net, GND = drift net, GTR – trammel net, OTB = bottom otter trawl and OTT = multirig bottom otter trawl) Taken from ICES (2021).

AU	Subarea	Métier L4	Vessel size	Ob- served DaS	Bycatch event/Da ^s			Number of individuals/bycatch event		
					mean	lower	upper	mean	lower	upper
NORTHSEA	27.3	GNS	small	1647	0.05	0.05	0.06	1.19	1.04	1.37
NORTHSEA	27.3	GNS	large	1782	0.40	0.27	0.55	1.16	0.99	1.35
NORTHSEA	27.3	GTR	small	82	0.05	0.05	0.06	1.19	1.04	1.37
NORTHSEA	27.3	GTR	large	0	0.40	0.27	0.55	1.16	0.99	1.35
NORTHSEA	27.3	OTB	All	21907	0.00	0.00	0.00	1.10	0.90	1.35
NORTHSEA	27.3	OTT	All	7486	0.00	0.00	0.00	1.10	0.90	1.35
NORTHSEA	27.4	GND	small	288	0.08	0.06	0.11	1.38	1.16	1.64
NORTHSEA	27.4	GND	large	3.91	0.02	0.01	0.05	1.33	1.15	0.46
NORTHSEA	27.4	GNS	small	1747	0.08	0.06	0.11	1.38	1.16	1.64
NORTHSEA	27.4	GNS	large	3.91	0.02	0.01	0.05	1.33	1.15	0.46
NORTHSEA	27.4	GTR	small	1073	0.08	0.06	0.11	1.38	1.16	1.64
NORTHSEA	27.4	GTR	large	3.91	0.02	0.01	0.05	1.33	1.15	0.46
NORTHSEA	27.4	OTB	All	50951	0.00	0.00	0.00	1.10	0.90	1.35
NORTHSEA	27.4	OTT	All	6392	0.00	0.00	0.00	1.10	0.90	1.35
NORTHSEA	27.7	GND	small	67	0.01	0.01	0.01	1.06	0.94	1.19
NORTHSEA	27.7	GND	large	0	0.03	0.02	0.05	1.18	0.94	1.49
NORTHSEA	27.7	GNS	small	4789	0.01	0.01	0.01	1.06	0.94	1.19
NORTHSEA	27.7	GNS	large	0	0.03	0.02	0.05	1.18	0.94	1.49
NORTHSEA	27.7	GTR	small	6068	0.01	0.01	0.01	1.06	0.94	1.19
NORTHSEA	27.7	GTR	large	322	0.03	0.02	0.05	1.18	0.94	1.49
NORTHSEA	27.7	OTB	All	16842	0.00	0.00	0.00	1.10	0.90	1.35
NORTHSEA	27.7	OTT	All	567	0.00	0.00	0.00	1.10	0.90	1.35

Table 4. Estimated bycatch of harbour porpoise in the Greater North Sea assessment unit and métier level 4 in 2019 and 2020. Numbers of individuals taken as bycatch are obtained by multiplying the average bycatch rates (animals caught per day-at-sea) by the annual fishing effort. Lower and upper values represent 95% confidence intervals. Adapted from ICES Advice (2021b). * Evidence of non-random sampling.

Métier level 4	Estimated bycatch rate 2015–2020 (95% CI)	Number of individuals taken as bycatch 2019 (95% CI)	Number of individuals taken as bycatch 2020 (95% CI)
Gill and drift nets*	0.240 (0.137–0.409)	5696 (3021–10391)	5327 (2845–9637)
Trammel nets*	0.247 (0.142–0.418)	690 (399–1178)	479 (277–821)
Bottom otter and multi-rig otter trawls	0.001 (0.0005–0.003)	145 (64–331)	123 (54–281)
All		10096 (5799–18198)	9299 (5374–16677)

Table 5: Summary of actual and potential pressures on the population.

Actual/Potential Threat	Cause or related activity	Evidence	Possible Impact	Priority for Action
Primary pressures				
Bycatch – lethal entanglement in fishing gears	Commercial and recreational static nets and trawls	Strong	Mortality	High
	Marine debris (including ghost nets)	Weak	Mortality and morbidity	Low
Serious/fatal injury (not bycatch)	Ship strikes from commercial and recreational vessels	Weak	Mortality or compromising injury	Low
	Collision with wet renewables	Moderate	Mortality or compromising injury	Low
Secondary pressures				
Mechanical destruction of habitat	Bottom trawls	Weak	Reduction in prey species	Low
	Infrastructure construction, oil and gas development			
	Gravel extraction			
Prey depletion	Overfishing	Moderate	Loss of body condition/reduced nutritional status, suppression of reproduction, mortality	Low (further evidence required)
	Habitat degradation due to pollution			
Chemical pollution	Atmospheric transportation, terrestrial industrial development, landfill, terrestrial run-off, harbours, ships, aquaculture, sewer discharges, aerial transport, oil spill	Strong	Immuno-suppression, increased disease risk, reproductive failure and dysfunction	Medium

Tertiary pressures				
Noise Disturbance	Fishing vessels, maritime traffic, recreational activities	Moderate	Displacement or injury	Medium
	Acoustic deterrent devices at fish farms, e.g. pingers			
	Military activities			
	Infrastructure construction, oil and gas development (including seismic),			
	Aggregate extraction			
Boat-based dolphin watching and other recreational activities		Moderate	Reduced foraging	Low
Environmental change	Further environmental changes are likely to affect marine conditions	Moderate	Change in distribution, and availability of prey and habitat	Medium
Cumulative impacts	The cumulative impact of pressures will increase risk to the population	Moderate	Reduced resilience to pressures due to combined impacts	Medium

4.2. Attributes of the population for monitoring, mitigation and research

To address the pressures summarised above, there is a requirement for monitoring, mitigation and/or research. For example, bycatch has been identified as the greatest anthropogenic pressure on this species. There remains a degree of uncertainty in the assessment of population bycatch rates due to ambiguities in recording fishing effort, biases and unrepresentative sampling by gear type, and a lack of statutory reporting from some major fishing nations (Advice 2016, ICES Advice 2021a, b). Other pressures in the region include marine pollution and underwater noise, with major knowledge gaps in the extent of their effects which hinder the provision of robust scientific assessments.

The attributes that have been identified as requiring monitoring, mitigation or research are listed in Table 6. Measures by which to assess the success of actions will be developed alongside each action by the Steering Group.

Table 6: Attributes for monitoring, mitigation and research.

Attribute	Relevant actions
Bycatch: High and medium risk fisheries and gear types, bycatch rates, effectiveness of mitigation measures including gear modifications	RES-01; RES-02; MIT-01; MON-01; RES-03; RES-04
Harbour porpoise health: Health and nutritional status, life history parameters and contaminant levels (and possible sources)	MON-02; MIT-02; MON-03; MON-04; RES-04
Noise pollution: Risks and impacts of underwater noise including renewable energy developments	MON-01; RES-03; MIT-02; MON-04
Cumulative impacts: Impact of activities in combination	MON-04; RES-02; RES-03; MON-02; MON-03;
Emerging pressures: Environmental change, pollutants of emerging concern	MON-01; RES-03; MON-02; MON-04
Conservation status: Population structure, demography and viability	RES-02; MON-01; RES-03; RES-04; MON-05

5. Conservation Status

Assessment of conservation status requires consideration of changes in range and abundance, as well as habitat preferences and availability. The assessment also requires an understanding of the main pressures and threats to the species that may impact longer term survival and also the population context against which the effectiveness of management of those pressures can be judged.

Because the range of harbour porpoise extends beyond the Greater North Sea, this element of the assessment is largely unchanging. Fluctuations in the population (i.e. abundance and trends) are therefore the more important determinant of conservation status, particularly where links can be made to anthropogenic activities that may cause declines. Changes in distribution within range can also be important. For example, by the 1940s the harbour porpoise had become rare in the southern North Sea and English Channel probably as a result of overfishing, bycatch and/or local changes in oceanographic conditions leading to changes in pelagic assemblages including dominant fish populations, with reoccupation only beginning in the 1990s (Reid et al. 2003, Hammond et al. 2013, Murphy et al. 2013, IAMMWG 2015, Evans 2020).

Table 7: Conservation status conclusions for harbour porpoise in the European Marine Atlantic biogeographic region (FRP – favourable reference population). (From Pinn et al., 2021).

Country	Assessment period		
	2001-2006	2007-2012	2013-2018
Belgium	Unfavourable bad	Unfavourable inadequate	Unknown
Denmark	Unfavourable bad	Favourable	Favourable
France	Unknown	Unfavourable bad	Unfavourable inadequate
Germany	Unfavourable inadequate	Unfavourable inadequate	Unfavourable inadequate
Ireland	Favourable	Favourable	Favourable
Netherlands	Unfavourable bad	Unfavourable inadequate	Favourable
Portugal	Unfavourable inadequate	Unfavourable inadequate	Unfavourable bad
Spain	Unknown	Unfavourable inadequate	Unfavourable inadequate
Sweden	Unfavourable bad	Unfavourable bad	Favourable
United Kingdom	Favourable	Favourable	Unknown
Overall conclusion for European Marine Atlantic region	Unfavourable-inadequate (80% of population in a favourable condition)	Favourable (89% of population in favourable condition)	Favourable (64% of population reported as favourable. UK assessment of unknown, covering 27% of population, was considered overly precautionary).

5.1. Critical Habitats

Through the Habitats Directive, EU Member States have a commitment to identify Special Areas of Conservation (SACs) for harbour porpoise. Article 4(1) notes that the designation of SACs for wide ranging aquatic species such as harbour porpoise '*will be proposed only where there is a clearly identifiable area representing the physical and biological factors essential to their life and reproduction*'.

Annex III of the Directive sets out general criteria for selecting SACs:

- *'Criterion a. Size and density of the population of the species present on the site in relation to the populations present within the national territory;*
- *Criterion b. Degree of conservation of the features of the habitat which are important for the species concerned and restoration possibilities;*
- *Criterion c. Degree of isolation of the population present on the site in relation to the natural range of the species; and*
- *Criterion d. Global (overall) assessment of the value of the site for the conservation of the species concerned.'*

By 2020, 232 sites had been designated in European Union waters for harbour porpoise (EEA, 2020, Figure 8). Although now no longer part of the EU, the UK has retained their designated SACs, which have also been listed as part of the Emerald Network under the Bern Convention.

The designation of SACs places specific duties on public authorities to manage activities they are responsible for in a way that avoids site deterioration and ensures protection of important species habitat. However, the value of such areas is severely diminished if the threats to the species are not tackled appropriately. Notably, for members of the EU, the management of fisheries has been delegated to the European Commission. As a result, whilst Member States' have a responsibility to manage their own SACs, they are unable to impose fisheries measures. Instead, if fisheries measures are required to achieve the conservation objectives of the SAC, they must be requested and implemented through the Common Fisheries Policy (CFP).

For the Greater North Sea, any fisheries measures are negotiated and agreed via the Scheveningen Group, which comprises the Fisheries Directors of the North Sea Member States. Once agreed, the proposed fisheries measures are then submitted to the Scientific, Technical and Economic Committee for Fisheries (STECF) for assessment. STECF either return the measures to the Scheveningen Group for revision or can advise that the measures are adopted and implemented through the CFP. Following over a decade of negotiations, in December 2022, the European Commission finally adopted fisheries measures banning the use of gillnets to protect harbour porpoise in six SACs in the North Sea. These are the German sites Sylt Outer Reef, Borkum Reef Ground, Dogger Bank and Eastern German Bight; and the Dutch sites Cleaver Bank and Frisian Front³.

Since leaving the EU, the UK is now in a unique position and is able to implement fisheries measures within its protected sites with immediate effect. The Trade and Cooperation Agreement⁴ between the UK and EU allows for this, provided that the measures are based on best available scientific evidence and that the same measures are applied to both UK and EU vessels. Initially the UK focus has been on the fisheries measures required to protect habitat features within its SAC network. It is expected that consideration will be given to the need for fisheries measures in the harbour porpoise SACs in 2024/5.

³ https://oceans-and-fisheries.ec.europa.eu/publications/c2022-8918_en

⁴

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/982648/TS_8.2021_UK_EU_EAEC_Trade_and_Cooperation_Agreement.pdf




Figure 8: Marine Atlantic regional network of Special Areas of Conservation for harbour porpoise to end of 2019. Sites with harbour porpoise as a qualifying feature (grades A–C) are shown with blue stripes and those where the species is a non-qualifying feature (grade D) in red. The Marine Atlantic biogeographic region is shown in pale blue (From Pinn et al., (2021)).

5.2. Dealing with inadequate data

While ideally, all conservation plans and associated management actions are based on full and adequate scientific data, there are occasions when the potential conservation consequences of waiting for confirmatory scientific evidence may mean that it is better to take action in the short term whilst collecting further evidence. This has become known as following the Precautionary Principle⁵. However, application of the precautionary principle must be carefully considered and adequately justified. One of the main challenges encountered in the process of developing the original version of the CMP (ASCOBANS, 2009) was the lack of data available on which to base some decisions. This issue still persists today.

The actions, therefore, include a number of research and monitoring actions which work towards obtaining the necessary information for the establishment of adequate scientifically-based management actions. These actions need to be given some priority to ensure management or mitigation is based on robust data and therefore likely to be effective.

⁵ <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=LEGISSUM%3AI32042>

6. Actions

6.1. Summary of actions

Below is a list of the identified actions, with an indication of priority and likely constraints of achieving each. Actions are categorised under Monitoring (MON); Mitigation (MIT) and Research (RES) codes.

Priority	Action	Code	Constraints
Essential	Identify the priority bycatch issues and relevant stakeholders	RES-01	Political will influenced by societal desire to support
Essential	Improve estimates of bycatch rates to support development of conservation strategy	RES-02	Metrics used to record fishing effort; ambiguous definitions for some gear types; insufficient funding to support the extent of monitoring needed for robust estimates
Essential	Implement and assess pinger and mitigation measures to reduce bycatch	MIT-01	Cooperation from fishing industry; enforcement measures
High	Implement a wide-scale surveillance programme to monitor trends in distribution and abundance in the Greater North Sea	MON-01	Commitment of funding
High	Improve understanding of causes of seasonal and annual variation in abundance and distribution, particularly in relation to human activities and environmental change, to facilitate consideration of the species within marine spatial plans	RES-03	Although this is one of the most surveyed regions in the North-east Atlantic, the spatial temporal coverage is still inadequate, thus there are difficulties in mapping some human activities/impacts
High	Monitoring of health and nutritional status, diet, life history parameters, and causes of mortality	MON-02	Commitment of funding; access to samples; development of suitable methods
Medium	Further our understanding of population structure	RES-04	Development of non-invasive sampling methods; discrimination ability of different techniques.

Priority	Action	Code	Constraints
Medium	Improve understanding of and develop mitigation for the risks of anthropogenic sound	MIT-02	Difficulty in attributing sound exposure to physical or behavioural consequences at both the individual and population level
Medium	Ensure screening and assessment of the occurrence and effects of hazardous substances	MON-03	Effective identification of emerging hazards; addressing impacts on harbour porpoises specifically
Low	Monitor for potential increases in anthropogenic activities that lead to incidences of death, injury or adverse health effects	MON-04	Availability and accessibility of information
Low	Monitor habitat quality, including protected sites, to ensure management is effective and that the ecological functions are maintained.	MON-05	Political will influenced by societal desire to support

6.2. Actions and Tasks

The actions are detailed below setting out the priority tasks and constraints to achieving the action objectives, and who is responsible. Monitoring actions identify key tasks in developing monitoring programmes for the species, similarly with Mitigation actions. Research actions identify tasks essential for providing adequate management advice. The tasks identified within each action will formulate the basis on which countries will report progress to ASCOBANS.

The North Sea CMP Steering Group (NSSG) will be responsible for developing detailed plans for tasks where required to coordinate implementation and identify a way forward. The NSSG will collate reports on the progress of implementation, effectiveness and issues encountered and report annually to the Advisory Committee on the progress of the CMP, identifying further implementation priorities and make appropriate recommendations. The reporting will be concise and efficient to reduce burden and maintain up to date information on application and progress of tasks.

Action RES-01: Identify the priority issues and relevant stakeholders

Priority: ESSENTIAL

Research action

Constraints: depends on political will, influenced by public support.

Description of action

Static net fisheries are recognised as being the greatest risk to harbour porpoise. There is a need to identify those of highest risk in terms of temporal and spatial extent, in order to effectively direct monitoring and mitigation effort. There is then opportunity to:

- prioritise mitigation measures, management and innovation to address ASCOBANS conservation objectives.
- improve understanding of the factors which influence bycatch levels; e.g. age, sex, time of day of capture, season, location, hydro-meteorological condition, associated prey species, gear specifications and usages etc.;
- facilitate implementation of the management framework procedure and indicators of bycatch developed by OSPAR to support collaborative approaches at an appropriate spatial scale.

Attention is needed to revise the current ASCOBANS conservation objectives to incorporate a timeframe for their achievement, and to take account of the long-term objective to drive anthropogenic removals towards zero mortality.

Engagement with other relevant stakeholders, including fishers and fisheries Regional Coordination Group North Atlantic, North Sea and Eastern Arctic (NANSEA), as well as scientists, NGOs and government managers, is required to reach common solutions and to fulfil conservation objectives.

Tasks

1. Identify and monitor medium-to-high-risk static net and other fisheries with a medium-to-high risk of harbour porpoise bycatch in order to ascertain more accurate assessments of bycatch rates to meet the agreed objectives of Resolution 3 MOP 3, Resolution 5 MOP 8 and Resolution 8.5 MOP9.
2. Implement the management framework procedure developed through OSPAR and progress development of suitable indicators of bycatch for the harbour porpoise with other fora, to meet the agreed objectives of Resolution 5, MOP 8 and Resolution 8.5 MOP9.
3. Facilitate the identification of factors influencing bycatch rates; including an assessment of temporal (seasonal) and spatial, gear characteristics, fishing practices and target/non-target species.
4. Facilitate research in order to assess evidence of bycatch selectivity of age-sex groups in different fishing operations (e.g. gears, target species, seasons), with the inclusion of those data within a population viability analysis.
5. Monitor causes of death in the population through strandings programmes for aiding assessments of spatio-temporal relationships and trends in bycatch, aiding implementation of the agreed objectives of Resolution 10, MOP8 and Resolution 8.10 MOP9 on strandings.
6. Represent ASCOBANS and the North Sea Plan at meetings of NANSEA and the North Sea Regional Advisory Council, as well as engagement with Parties' fisheries administrations.

Actors

CMP Coordinator/Steering Group, ASCOBANS-ACCOBAMS JBWG, national authorities, other stakeholders including OSPAR and scientists (e.g. ICES WGBYC), NANSEA, and North Sea Regional Advisory Council.

Action RES-02: Improve estimates of bycatch rates to support development of conservation strategy

Priority: ESSENTIAL

Research action

Constraints: Potential constraints are the current metrics used to record fishing effort, ambiguous definitions for some gear types, insufficient funding or inefficient use of available funding to support the extent of monitoring needed for robust estimates.

Description of action

Bycatch estimates across the Agreement area are hampered by low sampling effort and the difficulties in quantifying effort adequately due to the format of recorded information from relevant fisheries. Currently, effort is logged as days at sea rather than more accurate measures that take account of net dimensions and soak times (e.g. (ICES 2022)). Bycatch rates are determined from visual observers aboard a small fraction of active vessels, as well as some remote electronic monitoring (REM). Although EU Range States are requested by ICES to report bycatch rates on an annual basis, some do not, or data submissions are incomplete. Efforts are needed at international, national and regional levels to improve the level and frequency of provision of information. There still remains great uncertainty around all bycatch estimates for the harbour porpoise in the Greater North Sea.

Tasks

1. Ensure that existing regulations with respect to bycatch reduction measures are being effectively implemented and to collect data on their efficacy in reducing bycatch to meet the agreed objectives of Resolution 3, MOP 3, Resolution 5, MOP 8 and Resolution 8.5 MOP9.
2. Drive coordination of bycatch monitoring observer programmes across Parties and non-Party Range States, ensuring that monitoring programmes have been designed appropriately, with a sufficient level of monitoring to produce robust and unbiased estimates of bycatch with confidence intervals.
3. Increase reliability of fishing effort data particularly for small vessels (<12 m)⁶ and recreational fisheries and continue evaluating appropriate fishing effort metrics for calculating bycatch rates, supporting the wider work of ICES. This involves, working nationally (e.g. through work plans) and regionally (through Regional Coordination Groups) to improve quality and availability of fishing effort data (e.g. by region, gear type, net length, vessel size category, season, and country).
4. Support innovative monitoring methods, e.g. REM, particularly for use on smaller vessels (<12 m) where the placing of onboard observers is not feasible, and liaise with ICES WGBYC on how best these data should be collated and assessed as different monitoring methods will have different levels of uncertainty.
5. Support the development of OSPAR's M6 Biodiversity Common Indicator 'Marine Mammal Bycatch', which will aid EU Member States in meeting requirements of the MSFD as well as agreed objectives of ASCOBANS Resolution 3, MOP 3, Resolution 5, MOP 8 and Resolution 8.5 MOP9.

Actors

CMP Coordinator/Steering Group and ASCOBANS-ACCOBAMS JBWG, with support from Range States/Parties to ASCOBANS and ICES.

⁶ This is required by the Habitats Directive where bycatch from small vessels is thought to have a negative impact on conservations status. It is also required by the EU Data Collection Framework Regulation 2017/1004 (<https://datacollection.jrc.ec.europa.eu/legislation/current/obligations>) and the EU Implementing Regulation 2019/1241.

Action MIT-01: Implement and assess pinger and other mitigation measures to reduce bycatch

Priority: ESSENTIAL

Mitigation Measure Action

Constraints: Political will, socio-economic cost and willingness of industry.

Description of action

The use of pingers in certain static net fisheries was mandated in 2004 through EU Regulation 812/2004. This regulation has since been repealed and the requirements incorporated into EU Data Collection Framework Regulation 2017/1004⁷ and the EU Implementing Regulation 2019/1241⁸. Today the legislative emphasis is on international commitments for protected species, incorporating all fisheries that may have a negative impact. This also reflects the requirements of the EU Habitats Directive for monitoring of bycatch and implementing mitigation measures where there is a negative impact on conservation status. The UK requires the use of pingers in those fisheries where it was originally mandated. Any other vessel (e.g. all inshore vessels) is required to obtain a licence to use pingers. In contrast, all coastal gillnet vessels are required to use pingers in Norway to reduce bycatch.

Since their introduction, it has become clear that pingers are very effective in some fisheries but not in others (ICES WGBYC 2020, Lusseau et al. 2023). There is also a need to further understand the contradictory evidence on the possible effects of habituation and habitat exclusion in relation to pinger deployment. Given these concerns, the use of alternative gear types is often advocated (Leaper and Calderan 2018, Read 2021). However, due to the cost of switching gear, relicensing a vessel and learning to fish using a different technique, this approach is unviable for many smaller vessels (Ryan et al. 2022). A focus on gear adaptation has therefore been advocated by industry.

The ultimate aim for the development of any mitigation measure is to ensure universal acceptance by all stakeholders (and hence better implementation) of mitigation measures to reduce harbour porpoise bycatch.

Tasks

1. Implement mitigation measures that have shown to produce a significant bycatch reduction and that are appropriate to the nature of the vessels and their size, with subsequent monitoring to ensure effectiveness and the ongoing need to meet the agreed objectives of Resolution 5, MOP 8 and Resolution 8.5 MOP9. It may be necessary to undertake an Environmental Risk Assessment for the implementation of pingers *en masse*.
2. Collaborate with the industry to develop and test mitigation measures (including modifications to fishing gear and fishing practices; pinger-related technology and deployment (e.g. interactive pingers, less pingers per length of net), and alternative porpoise alerting passive and active devices) and develop a framework for the critical evaluation of pinger, gear modification and other mitigation measures to identify effectiveness in the reduction of bycatch to meet the agreed objectives of Resolution 5, MOP 8 and Resolution 8.5 MOP9.
3. Support research evaluating the behaviour of harbour porpoises around fishing gear, especially static nets, including their sensory capabilities and auditory health, for a better understanding of factors leading to bycatch.
4. Prevent, retrieve, and recycle derelict (“ghost”) fishing gear, with focus on high-density areas of harbour porpoises as agreed by Resolution 9.3 MOP9. This will require authorities to provide appropriate facilities to ensure gear is recycled and to prevent disposal of at sea.

Actors

CMP Coordinator/Steering Group and ASCOBANS-ACCOBAMS JBWG, with support from Range States and Parties to ASCOBANS, fisheries authorities and scientists.

⁷ <https://datacollection.jrc.ec.europa.eu/legislation/current/obligations>

⁸ <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32019R1241>

Action MON-01: Implement a wide-scale surveillance programme to monitor trends in distribution and abundance in the Greater North Sea

Priority: HIGH

Monitoring Action

Constraints: Availability of funding which may be driven, in part, by political will and support for the CMP.

Description of action

Information on trends in abundance and distribution is essential for the contextualisation of the majority of the actions associated with this CMP. Without such monitoring, it will be impossible to evaluate the success or otherwise of the CMP and to determine whether any modifications are required.

The fundamental basis for determining changes in harbour porpoise conservation status within the Greater North Sea is a programme of regular wide-scale standardised surveys. Given the cost, the term 'regular' would need to be identified based on sufficiency for reporting trends. Recent work has deemed that SCANS-types surveys should be undertaken at a six-year frequency. However, these surveys provide 'snapshots' of the abundance and distribution within the area surveyed, typically being carried out over a one-month period during the summer. Given the temporal limitations, complimentary coordinated regional data collection is also required to ascertain long-term and seasonal changes in distribution at a North Sea-wide scale and the examination of potential explanations for any observed changes.

Tasks

1. Encourage Parties and non-Party Range States to collaborate and fund regular systematic North Sea-wide and regional surveys to establish trends in abundance and distribution, and undertake density surface modelling, to meet the agreed objectives of Resolution 7, MOP4 and Resolution 7, MOP5.
2. Encourage Parties and non-Party Range States active participation with the ICES Working Group on the Joint Cetacean Data Programme (WGJCDP) which has developed a mechanism for collation of all relevant, standardised data at a relevant spatial scale, collected through ship-based and aerial methodologies, and aims to develop analyses and data products in line with identified priorities across the cetacean research and policy community. Such work would enable seasonal trends to be evaluated to meet the agreed objectives of Resolution 7, MOP4.
3. Ensure Parties and non-Party Range States that the outputs of this action provide a suitable mechanism to enhance transboundary reporting of conservation status and good environmental status, as well as contributing to the assessment of OSPAR's M4 Biodiversity Common Indicator 'Abundance and Distribution of Marine Mammals', evaluating temporal trends and any further re-distribution of individuals within the Greater North Sea.

Actors

CMP Coordinator/Steering Group with support from Range States/Parties to ASCOBANS, OSPAR, scientists and managers especially those involved in the monitoring component of SCANS and other surveillance work.

Action RES-03: Improve understanding of causes of seasonal and annual variation in abundance and distribution, particularly in relation to human activities and environmental change, to facilitate the consideration of the species within marine spatial plans

Priority: HIGH

Research action

Constraints: Although this is one of the most surveyed regions in the North-east Atlantic, the spatial temporal coverage is still inadequate, thus there are difficulties in mapping some human activities/impacts.

Description of action

A wide variety of anthropogenic activities occur in the greater North Sea region that may potentially affect harbour porpoises. It is necessary to be able to determine the occurrence and temporal/geographical distribution of these and any changes over time to be able to:

- compare these with the distribution of the animals to determine potential problem areas;
- to have baseline information to compare if changes in harbour porpoise abundance and distribution are observed via MON-01 (Implement a wide-scale surveillance programme to monitor trends in distribution and abundance in the greater North Sea).

Analyses should investigate relationships between the distribution and trends regarding relevant human activities (linking to Action RES-01: Identify the priority bycatch issues and relevant stakeholders and Action MIT-02: Improve understanding of and develop mitigation for the risks of anthropogenic sound) and climate-related indicators. Consideration of indirect impacts of environmental change (e.g. availability and re-distribution of preferred prey) should be considered where possible. This may be of particular importance for harbour porpoise SACs that likely encompass important foraging areas (see MON-05 Monitor habitat quality, including protected sites, to ensure management is effective and that the ecological functions are maintained).

Tasks

1. Continued collection and collation of appropriate standardised data on anthropogenic activities with the aim of supporting implementation of the MSFD and assessment of Good Environmental Status through OSPAR.
2. Complete fine-scale seasonal risk assessment/risk mapping of relevant human activities and harbour porpoise distribution to meet the agreed objectives of Resolution 7, MOP4, Resolution 7, MOP5 and Resolution 5, MOP8.
3. Collate and monitor data on important prey species of harbour porpoises to identify spatial areas of concern for fisheries management measures to meet the agreed objectives of Resolution 7, MOP4, Resolution 7, MOP 5 and Resolution 9.4 MOP9.
4. Through collaboration with other ASCOBANS working groups, such as the Resource Depletion Working Group, regularly review of evidence for potential impacts of environmental change on harbour porpoises to inform on appropriate mitigation measures.

Actors

Range States/Parties to ASCOBANS; scientists and managers especially those involved in the monitoring component of SCANS, Data collectors, fisheries authorities, OSPAR, ICES, policymakers, CMP Coordinator/Steering Group, contractors.

Action MON-02: Monitoring of health and nutritional status, diet, life history parameters and causes of mortality.

Priority: HIGH

Monitoring Action

Constraints: Funding; access to sufficient samples across the region.

Description of action

Our knowledge of the qualitative and quantitative effects of a range of human activities on harbour porpoises is incomplete. This impacts our ability to determine their conservation status and implement relevant good environmental status indicators.

This action is designed to improve this by collecting and reviewing information on causes of mortality, health and nutritional status, diet, as well as life history parameters. Types of data also required for population dynamics modelling.

Information on diet and various health and life history parameters has historically been obtained from dead animals that have stranded, or in some cases been recovered as bycatch, which remains the primary source of these data.

Tasks

1. Fund national stranding and bycatch observer programmes and undertake full necropsies on a representative sample of carcasses (sex, age), for assessing cause of death, health status, diet, and life history parameters, to meet the agreed objectives of Resolution 10, MOP8 and Resolution 8.10 MOP9.
2. Ensure implementation of the joint ASCOBANS and ACCOBAMS 'Best Practice on Cetacean Post-mortem Investigation and Tissue Sampling'⁹ to achieve standardised, comparable datasets.
3. Encourage collaboration between stranding networks in the event of an unusual mortality event to identify potential causes of death, as well as encouraging collaborative research on the extent and potential reasons for grey seal predation, starvation/emaciation and acoustic trauma observed in harbour porpoises.
4. Support strandings programmes to enable the analysis of diet, including tissue samples for fatty acids, stable isotope, stomach contents, and prey DNA analysis.
5. Support North Sea-wide monitoring of life history parameters through the collection and analysis of teeth and gonadal samples from stranded and bycaught animals, to assess evidence of temporal changes in life history parameters and explore links to anthropogenic drivers.
6. Support expansion of drift prediction modelling capabilities for determining the origin of stranded harbour porpoises, e.g. MOHY (Peltier et al. 2013, Peltier et al. 2018) to identify potential bycatch high risk areas/seasons.
7. Support the development of a biodiversity 'population condition' indicator for the region.

Actors

Range States, EC, International Whaling Commission Scientific Committee, ASCOBANS, CMP Coordinator/SG, other stakeholders including OSPAR, scientists and strandings programme coordinators.

⁹ https://www.ascobans.org/sites/default/files/document/ascobans_mop9_doc6.2.5b_best-practice-cetacean-post-mortem-investigation_pdf

Action RES-04 Further our understanding of population structure

Priority: MEDIUM

Research action

Constraints: Potential constraints are the discrimination ability of different techniques, practicalities of introducing a well-designed sampling procedure, and development of acceptable non-invasive methods to collect the appropriate information.

Description of action

Information on population structure may be obtained by a variety of means, including, amongst others, DNA analysis (mtDNA, microsatellite, MHC (Major Histocompatibility Complex) and SNP (single nucleotide polymorphism) markers, whole genomic studies by new generation technologies), morphometric studies, stable isotope signatures, fatty acid profiles, and comparisons of life history parameters.

Each is characterised by having different powers of discrimination over different time scales. Traditionally, most information on the population has come opportunistically from strandings, though bycaught animals have been extensively sampled through European observer programmes. Strandings data offer valuable insight, however, have limitations. Therefore, methods to reduce these limitations (e.g. improved drift modelling) and methods of collecting more representative samples should be explored. Combining relevant approaches, such as population genetics, ecological tracers (e.g. contaminants, stable isotopes), and trends in life-history parameters, would provide a comprehensive picture of the multifarious dimensions of the ecology and evolution of harbour porpoises in the region (Murphy et al. 2019).

Tasks

1. To further research relevant to ASCOBANS objectives on population structure of the harbour porpoise, to improve conservation measures, as agreed in Resolution 7 MOP5.
2. Actively support and encourage development of suitable techniques for discriminating population structure, as agreed in Resolution 7, MOP5.
3. To identify funding and develop a programme of research to further elucidate the population structure of harbour porpoises in the region. Strategic sampling approaches (i.e. temporal and spatial) and statistical power analysis should be undertaken to determine level of sampling required to detect appropriate units to conserve.
4. Facilitate the provision of dead bycaught animals for population structure assessment and other appropriate studies. This may require repeal of national legislation or the issuing of licenses to facilitate landing of bycaught harbour porpoise for research.

Actors

Range States, CMP Coordinator/SG, other stakeholders including scientists, fisheries authorities and strandings programme coordinators.

Action MIT-02: Improve understanding of and develop mitigation for the risks of anthropogenic sound

Priority: MEDIUM

Mitigation Measure Action

Constraints: Difficulty in attributing sound exposure to physical or behavioural consequences at both the individual and population level.

Description of action

A wide variety of anthropogenic activities introduce sound into the marine environment, e.g. vessels, construction and operation of windfarms, general construction works, hydrocarbon exploration, military activities including removal of munitions, pingers, acoustic harassment devices. However, the actual or potential effects of such sounds on harbour porpoises in the short-term or long-term has not been fully quantified. Individual based modelling frameworks, such as iPCoD (Mortensen and Thomsen 2019) and DEPONS (Nabe-Nielsen et al. 2018), have been developed that further our understanding of the impacts of noise on vital parameters, though they require accurate and relevant input data.

GES indicators for noise have been developed which require substantial monitoring and reporting of noise activities. These are, however, limited to loud impulsive sounds (e.g. pile driving and underwater explosions), and continuous noise (e.g. shipping traffic). Through the JOMOPANS project, soundscape maps of ambient noise are being developed as a GES Tool to enable marine managers to quantitatively and graphically assess the risk of impacts on indicator species in the North Sea¹⁰.

Tasks

1. Parties and non-Party Range States should introduce precautionary guidance on measures and procedures for all activities surrounding the development of renewable energy production and other noise-producing industry to minimise risks to populations and mitigate possible effects following current best practice as agreed in Resolution 2, MOP 6 and Resolution 8.11 MOP9.
2. Parties to make every effort to mitigate the effects of activities involving explosions of munitions, as agreed in Resolution 8 MOP8.
3. Parties and non-Party Range States should coordinate and support research on the effects of underwater noise on harbour porpoises, including further development of individual based modelling frameworks, to meet the agreed objectives of Resolution 4, MOP 5, Resolution 2, MOP 6, Resolutions 6, 8 and 9, MOP 8 and Resolution 8.11 MOP9.
4. Annually monitor and assess knowledge of the effects of anthropogenic sound through review of literature, including acoustic capabilities of harbour porpoises, behavioural responses of harbour porpoises and the effectiveness of mitigation technologies as agreed in Resolution 2, MOP6 and Resolution 6, MOP8.
5. Support the work of MSFD Common Implementation Strategy Technical Group on Underwater Noise (TG-NOISE), and for Parties to implement agreed thresholds as they are developed (e.g. common methodology for assessment of impulsive underwater noise and continuous noise).

Actors

TG-NOISE, CMP Coordinator/Steering Group, national authorities, other stakeholders including OSPAR and scientists.

¹⁰ https://ec.europa.eu/environment/marine/pdf/NOISE%20Overview%20of%20main%20European-funded%20projects%20March_2022_Final.pdf

Action MON-03: Ensure screening and assessment of the occurrence and effects of hazardous substances

Priority: MEDIUM

Monitoring Action

Constraints: Identifying new products as hazardous; assessing impacts that apply specifically to the harbour porpoise.

Description of action

Programmes currently exist in the Agreement Area that monitor a suite of hazardous chemicals. However, the impacts that some of these may have specifically on harbour porpoises has not been fully assessed. In addition, assessment of levels of new emerging contaminants of concern on the EU watchlist for emerging pollutants is ongoing (Commission Implementing Decision (EU) 2015/495). This is particularly relevant for those pollutants identified as endocrine disrupting chemicals, which are known to effect health status (Law et al. 2012, Murphy et al. 2015, Jepson et al. 2016, Murphy et al. 2018, Williams et al. 2020, van den Heuvel-Greve et al. 2021, Williams et al. 2023).

Tasks

1. Continue collecting, archiving and analysing representative samples of porpoise tissues for relevant contaminants, with associated data on cause of death, health and nutritional status, and life history (linked to Action MON-02: Monitoring of health and nutritional status, diet, life history parameters, and causes of mortality; and RES-04: Further our understanding of population structure). Further assessment of confounding factors such as age, body condition, reproductive activity, and health status on individual pollutant loads should be undertaken.
2. Continue to monitor and assess emerging chemical pollutants and marine litter (including macro-, micro- and nanoplastics) in harbour porpoises through review of literature to progress agreed objectives of Resolution 4, MOP 7, Resolution 7, MOP5, Resolution 7, MOP8, and Resolutions 9.3 and 9.4, MOP9. Such work should devise a North Sea-based risk list of priority pollutants for monitoring in the species.
3. Monitor effects from exposure to legacy pollutants on immune, endocrine and reproductive functions in harbour porpoises against agreed toxicity thresholds, through continued analysis of strandings data to meet agreed objectives of Resolution 7, MOP8.
4. Encourage Parties and non-Party Range States to further develop thresholds to be employed for contaminants of concern, including the continued development of dose-response relationships between contaminants and physiological (reproductive and immunological) impacts for the harbour porpoise.
5. Encourage Parties and non-Party Range States to work through OSPAR and other relevant fora to aid the development of a marine mammal persistent chemical contaminants indicator of GES to meet Criteria D8C2 in order to ascertain that the health of the species is not adversely affected due to contaminants, including cumulative and synergetic effects. For such work, collection of a sufficient number of stranded and/or bycaught harbour porpoises is required to assess trends and status of persistent chemicals in the harbour porpoise assessment unit (linked to RES-04 Further our understanding of population structure).

Actors

Range states, other stakeholders including scientists, CMP Coordinator/Steering Group.

Action MON-04: Monitor for potential increases in anthropogenic activities that lead to incidences of death, injury or adverse health effects including cumulative effects.

Priority: LOW

Monitoring Action

Constraints: Availability of and access to the necessary information.

Description of action

Where current exposure of some pressures may be viewed as sustainable with regards to the harbour porpoise in the Greater North Sea, increases in exposure of either a single pressure, or cumulative increases, may have a negative impact and requires monitoring to enable early detection of risk, and subsequent development of management. A number of human activities known to have negative impacts upon marine mammals can be monitored from information gathered as part of other surveillance and monitoring programmes and, therefore, a strategic approach to data collection should be explored. Frameworks are being developed by other fora, such as OSPARs cumulative effects assessment¹¹, and assistance in such work is encouraged.

Tasks

1. Encourage Parties and Range States to continue to give their full support to the activities related to applying an ecosystem approach to the management of human activities under the frameworks of OSPAR, HELCOM, the European Union and the Convention in Biological Diversity as agreed in Resolution 9, MOP8 and Resolutions 8.11 and 9.3 MOP9.
2. As part of the annual reporting for the CMP, collect and review information to monitor changes in exposure to key anthropogenic pressures, and the effects arising from such, to support OSPAR's cumulative effects assessment work.
3. Requests that Parties and Range States ensure that cross-sectoral and transboundary consultations take place as early as the planning stage of activities in marine areas (marine spatial planning) with the aim of identifying potential impacts and minimising or mitigating such impacts effectively as agreed in Resolutions 6 and 9, MOP8, particularly where such work occurs within or adjacent to protected sites of the harbour porpoise (see Action MON-05: Monitor habitat quality, including protected sites, to ensure ecological functions are maintained).
4. Identify emerging pressures (e.g. wet renewables and ecotourism) and ensure monitoring is in place to establish risk. These emerging pressures need to be considered in the context of those already existing, and to take impacts into account cumulatively.

Actors

Range States national authorities, OSPAR, International Maritime Organisation (IMO), International Whaling Commission (IWC), CMP Coordinator/Steering Group.

¹¹ <https://www.ospar.org/news/cumulative-effects-assessment>

Action MON-05: Monitor habitat quality, including protected sites, to ensure management is effective and that the ecological functions are maintained.

Priority: LOW

Monitoring Action

Constraints: Depends on political will.

Description of action

The designation of Special Areas of Conservation (SACs) is required for the harbour porpoise by Article 4 of the Habitats Directive. Although no longer part of the EU, the UK have still retained their designated SACs, which have also been listed as part of the emerald network under the Bern Convention.

SACs aim to safeguard the species at critical locations for their lifecycle, whether they are used for feeding, breeding, resting or other activities (although details of their behaviours in these areas are not well understood). Because harbour porpoise are highly mobile, SACs have an important role in safeguarding the inherent ecological conditions required.

The SAC network within the North Sea is considered to be largely complete. The focus is now on implementing appropriate management of these sites in order to achieve their conservation objectives. It is also recognized that in order for these sites to be effective, it will be necessary to implement management and mitigation of anthropogenic pressures outwith the SACs.

Tasks

1. As part of the annual reporting for the CMP, collect and review information on habitat quality and protected area condition within the Greater North Sea.
2. Review conservation objectives and the implementation of management measures for SACs, assessing whether the conservation objectives are fit for purpose and that the management is effective.
3. Collect and review information on anthropogenic activities within and adjacent to SACs, and whether they have a significant impact on harbor porpoises at those sites (Action RES-03: Improve understanding of causes of seasonal and annual variation in abundance and distribution, particularly in relation to human activities and environmental change, to facilitate the consideration of the species within marine spatial plans).
4. Encourage Parties and Range States to identify the location of any further suitable sites for the establishment of protected areas, and to implement appropriate management actions in these areas on their own or in the context of other intergovernmental bodies to ensure the protection of harbour porpoise as agreed in Resolution 7, MOP5.

Actors

Coordinator/Steering Group, national authorities, other stakeholders including OSPAR and scientists

7. Public awareness and capacity building

This CMP has been developed to collate knowledge and information on the species and develop a set of relevant actions to implement to conserve the species with an aim to maintain and, where necessary, restore the North Sea management unit to favourable conservation status.

Wider awareness of both the pressures and the activities which cause them, and also any successes of the plan, will support achievement of the aims. Education and awareness may also contribute to better reporting of sightings and impacts, leading to better data for decision making.

There is the capacity for misinformation through media following events such as strandings; bycatch discard and other impacts such as vessel strikes. The outreach proposed for this plan could be effectively undertaken by better use of the media, including the internet (e.g. through ASCOBANS and Range State webpages), and activities such as public lectures and education programmes. It is important to continue communication particularly with stakeholders who have an impact on the species (e.g. through activities such as fishing, and renewable development, etc.) to maintain communication channels and support action of relevant tasks, as well as work with other interested parties to publicise the work ongoing to conserve the species.

7.1 Public awareness tasks

1. All key milestones (e.g. timetables for actions; assessment of progress against objectives etc.) to be publicised through ASCOBANS and Range State media outlets in a coordinated manner agreed through the SG.
2. ASCOBANS webpages to host key documents and updates, to be publicised by SG members.
3. Presentation of the progress at relevant events and conferences.
4. Identification and publication of papers through journals and list servers/webpages to publicise lessons learned and successes.
5. Wider circulation of articles and news items through the media/social media to support the dissemination of factual information to the wider public.
6. Coordination with relevant NGO's with an interest in harbour porpoise, to join up approaches for public information campaigns.

Annex 1: International conventions and agreements

In the NE Atlantic, harbour porpoise are incorporated into a wide variety of legislation including national, European and international statutes and conventions, all with aims to protect, conserve, manage and study the species. In addition, there is other international legislation aimed at specific industries.

Full Title	Acronym/shorthand
1.1 United Nations Convention on the Law of the Sea	UNCLOS
1.2 Convention on Biological Diversity	CBD
1.3 Convention on International Trade in Endangered Species of Wild Fauna and Flora	CITES
1.4 The Convention on the Conservation of Migratory Species of Wild Animals & the Agreement on the Conservation of Small Cetaceans of the Baltic, NE Atlantic, Irish and North Seas	CMS & ASCOBANS
1.5 Convention on the Protection of the Marine Environment of the NE Atlantic	OSPAR
1.6 The Bern Convention	
1.7 European Directive of Natural Habitats and Wild Fauna and Flora (92/43/EEC)	Habitats Directive
1.8 International Convention for the Regulation of Whaling	IWC
1.9 Common Fisheries Policy	CFP
1.10 Marine Strategy Framework Directive	MSFD
1.11 Environmental Impact Assessment Directive	EIA
1.12 Strategic Environmental Assessment Directive	SEA

1.1. United Nations Convention on the Law of the Sea (UNCLOS)

UNCLOS governs all aspects of ocean space: Specifically, the convention states that contracting parties “shall cooperate with a view to the conservation of marine mammals and in the case of cetaceans shall in particular work through the appropriate international organisations for their conservation, management and study” and that signatories must take measures “necessary to protect and preserve rare or fragile ecosystems as well as the habitat of depleted, threatened or endangered species and other forms of marine life” (United Nations, 2001).

1.2. Convention on Biological Diversity (CBD)

The aim of CBD is conservation of biological diversity, the sustainable use of the components of biological diversity and a fair and equitable sharing of the benefits arising out of the utilization of genetic resources. Kunming-Montreal Global Biodiversity Framework, adopted in 2022, aims to halt and reverse biodiversity loss, implementing the three aims of CBD in a balanced manner. The framework also contributes to the achievement of the 2030 Agenda for Sustainable Development. The framework uses the theory of change, which recognizes that urgent policy action is required globally, regionally and nationally to achieve sustainable development so that the drivers of undesirable change that have exacerbated biodiversity loss will be reduced and/or reversed to allow for the recovery of all ecosystems and to achieve the Convention’s Vision of Living in Harmony with Nature by 2050.

1.3. Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)

CITES aims to regulate international trade in species that are endangered or may become endangered if their exploitation is not controlled (CITES, 2012). CITES is implemented within Europe through two EC regulations (338/97 and 865/06 as amended). Species covered under CITES are listed in three appendices, with harbour porpoise listed in Appendix 2. This means that trade in the species is permitted as long as the authorities have ascertained that it will not be detrimental to the survival of the species; that the specimen was not obtained in contravention of the laws of that state for the protection of fauna and flora; and that any living specimen will be so prepared and shipped that it minimizes the risk of injury, damage to health or cruel treatment.

1.4. The Convention on the Conservation of Migratory Species of Wild Animals (CMS) and the Agreement on the conservation of small cetaceans of the Baltic, NE Atlantic, Irish and North Seas (ASCOBANS)

The Convention on Migratory Species (CMS), or Bonn Convention, sets out general provisions for the protection and conservation of certain migratory marine mammals (CMS Secretariat, 2012). Harbour porpoise in the North Sea are listed in Appendix II. Appendix II includes species that have an unfavourable conservation status and that require international agreements for their conservation and management, as well as those that have a conservation status that would significantly benefit from the international cooperation that could be achieved by an international agreement.

One such agreement is the Agreement on the Conservation of Small Cetaceans in the Baltic, NE Atlantic, Irish and North Seas (ASCOBANS) and another the Agreement on the Conservation of Cetaceans of the Black Sea, Mediterranean Sea and Contiguous Atlantic Area (ACCOBAMS).

1.5. Convention on the Protection of the Marine Environment of the NE Atlantic (OSPAR)

The OSPAR Convention (replacing the Oslo and Paris Conventions) is the mechanism by which 15 governments of the coastal states of NW Europe, together with the European Commission, cooperate to protect the marine environment of the NE Atlantic with a particular focus on marine pollution, as well as providing for the conservation and protection of habitats and species.

Article 2(1)(a) states “the Contracting Parties shall, in accordance with the provisions of the Convention, take all possible steps to prevent and eliminate pollution and shall take the necessary measures to protect the maritime area against the adverse effects of human activities so as to safeguard human health and to conserve marine ecosystems and, when practicable, restore marine areas which have been adversely affected”.

Harbour porpoise are listed by OSPAR as a threatened and declining species, with a focus on tackling bycatch.

OSPAR is also providing the oversight for the regional sea assessments of Good Environmental Status required for the Greater North Sea and the Celtic Seas by the Marine Strategy Framework Directive (see section 1.10). These assessments incorporate GES indicators covering trends in harbour porpoise distribution and abundance (M4 Biodiversity Indicator), as well as bycatch (M6 Biodiversity Indicator) and pollution (Candidate Indicator Trends and Status of persistent chemicals in marine mammals). Full, updated indicator assessments will be published within OSPAR's Quality Status Report in 2023.

1.6. The Bern Convention

The Convention on the Conservation of European Wildlife and Natural Habitats (or the Bern Convention) covers most of the natural heritage of the European continent and extends to some states of Africa (European Union 1979). Harbour porpoise in the North Sea are listed in Appendix 2 ‘strictly Protected Fauna Species’, for which the following activities are prohibited:

- all forms of deliberate capture and keeping and deliberate killing;
- the deliberate damage to or destruction of breeding or resting sites;
- the deliberate disturbance of wild fauna, particularly during the period of breeding, rearing and hibernation, insofar as disturbance would be significant in relation to the objectives of this Convention;
- the possession of and internal trade in these animals, alive or dead, including stuffed animals and any readily recognisable part or derivative thereof, where this would contribute to the effectiveness of the provisions of this article.

There is also a requirement for contracting parties to coordinate “efforts for the protection of the migratory species specified in Appendices II and III whose range extends into their territories”. For Member States of the European Union, the provisions of the Bern Convention are largely taken up in the 1992 Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC), otherwise known as the ‘Habitats Directive’.

1.7. European Directive of Natural Habitats and Wild Fauna and Flora (92/43/EEC) (commonly known as the Habitats Directive) 1992

The Habitats Directive transposes the Bern Convention in EU law. Harbour porpoise are listed in Annex IV of the Habitats Directive as ‘Animal and Plant Species of Community Interest in Need of Strict Protection’. Article 11 requires Member States to monitor the conservation status of the habitats and species listed in the annexes; Article 17 requires an assessment of conservation status to be sent to the European Commission every 6 years. In

the Directive, conservation status is defined as “the sum of the influences acting on the species that may affect the long-term distribution and abundance of its populations”. Conservation status can be considered favourable if:

- population dynamics data indicate that the species is maintaining itself on a long-term basis as a viable component of its natural habitats;
- the natural range of the species is neither being reduced nor is likely to be reduced in the foreseeable future, and;
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Under Article 12, Member States shall take the requisite measures to establish a system of strict protection for the animal species listed in Annex IV(a) in their natural range, prohibiting: (a) all forms of deliberate capture or killing of specimens of these species in the wild (i.e. bycatch); (b) deliberate disturbance of these species, particularly during the period of breeding, rearing, hibernation and migration; and (d) deterioration or destruction of breeding sites or resting places. Member States are required to undertake further research or introduce conservation measures to ensure that incidental capture and killing does not have a significant negative impact on the species concerned. This is specifically relevant for harbour porpoise.

Under Articles 3 and 4, Member States’ contribute to an ecologically coherent network of protected areas known as Special Areas of Conservation (SACs) for those species listed in Annex II if suitable sites can be identified. This includes harbour porpoise.

Annex III of the Directive sets out general criteria for selecting SACs:

- ‘Criterion a. Size and density of the population of the species present on the site in relation to the populations present within the national territory;
- Criterion b. Degree of conservation of the features of the habitat which are important for the species concerned and restoration possibilities;
- Criterion c. Degree of isolation of the population present on the site in relation to the natural range of the species; and
- Criterion d. Global (overall) assessment of the value of the site for the conservation of the species concerned’.

Since the introduction of the Habitats Directive, Member States have had difficulties identifying suitable SACs for harbour porpoise, particularly in meeting the criterion covering the size and density of the population largely due to the mobility of the species.

1.8. International Convention for the Regulation of Whaling 1946

The International Whaling Commission (IWC) was set up under the International Convention for the Regulation of Whaling, which was signed in Washington, D.C., in December 1946 (IWC, 2012). The purpose of the convention is to “provide for the proper conservation of whale stocks and thus make possible the orderly development of the whaling industry”. Each year, the IWC Scientific Committee, through its Sub-- Committee on Small Cetaceans, identifies priority species/regions for consideration by a review. Topics considered include distribution, stock structure, abundance, seasonal movements, life history, ecology, and directed and incidental takes.

1.9. EU Common Fisheries Policy (CFP)

One of the objectives of Regulation EU 1380/2013 of the European Parliament and of the Council on the Common Fisheries Policy (CFP) is that the CFP shall implement the ecosystem-based approach to minimize negative impacts of fishing activities on the marine ecosystem. Such requirements are detailed in the Technical Measures Regulation (EU) 2019/1241, and

Article 3, Paragraph 2(b) notes '*ensure that incidental catches of sensitive marine species, including those listed under Directives 92/43/EEC and 2009/147/EC, that are a result of fishing, are minimised and where possible eliminated so that they do not represent a threat to the conservation status of these species*'. For this purpose, conservation measures such as modifications or additional devices to reduce incidental capture of endangered, threatened and protected species, or limitations on the use of certain fishing gears, shall be adopted. Also, highly relevant is the request that Member States should collect data on fleets and their fishing activities under the data collection framework (DCF) to support the CFP. Member States should manage the collected fisheries data and make them available to end-users and other interested parties. These data include biological, environmental, technical and socio-economic aspects, for example data on the impact of fisheries on biological resources and the marine ecosystem.

In December 2022, the European Commission adopted fisheries measures to protect harbour porpoise in six SACs in the North Sea. These are the German sites Sylt Outer Reef, Borkum Reef Ground, Dogger Bank and Eastern German Bight; and the Dutch sites Cleaver Bank and Frisian Front¹².

1.10. EU Marine Strategy Framework Directive (MSFD) 2008

The Marine Strategy Framework Directive (MSFD, Directive 2008/56/EC) requires Member States of the European Union to develop marine strategies that apply an ecosystem-based approach to the management of human activities while enabling a sustainable use of marine goods and services. Priority should be given to achieving or maintaining good environmental status in the community's marine environment, continuing its protection and preservation, and preventing subsequent deterioration¹³. To determine Good Environmental Status (GES), 11 qualitative descriptors have been selected. In 2017, OSPAR published its intermediate assessment for the 11 Descriptors which included harbour porpoise in Biodiversity Indicators M4 Cetacean Abundance and Distribution and M6 Marine Mammal Bycatch. The first EU-wide limit for underwater noise have been developed by the MSFD Technical Group on Underwater Noise. The threshold limit clarifies that to be in a "tolerable" status, no more than 20% of a given marine area can be exposed to continuous underwater noise over a year; and no more than 20% of a marine habitat can be exposed to impulsive noise over a given day, and no more than 10% over a year¹⁴.

1.11. EEU Environmental Impact Assessment (EIA) Directive (85/337/EEC) 1985

The EIA Directive (85/337/EEC) calls for assessment of the impacts on the environment of certain public and private projects which are defined in Annexes I and II of the Directive. A mandatory EIA is required of all projects listed under Annex I, which are considered to have significant effects on the environment. Projects listed under Annex II are at the discretion of Member States and subject to consideration by the national authorities as to whether an EIA is required, taking criteria detailed in Annex III into account. The majority of projects that may impact common dolphins, such as offshore renewable development, are listed under Annex II.

¹² https://oceans-and-fisheries.ec.europa.eu/news/fisheries-and-nature-conservation-increased-protection-natura-2000-sites-north-sea-2022-12-08_en

¹³ https://ec.europa.eu/environment/marine/eu-coast-and-marine-policy/marine-strategy-framework-directive/index_en.htm

¹⁴ https://environment.ec.europa.eu/news/zero-pollution-and-biodiversity-first-ever-eu-wide-limits-underwater-noise-2022-11-29_en

1.12. EU Strategic Environmental Assessment (SEA) Directive (2001/42/EC) 2003

The SEA Directive calls for an environmental report in which the likely significant effects on the environment and the reasonable alternatives of the proposed plan or programme are identified. The public and the environmental authorities are informed and consulted on the draft plan or programme and the environmental report prepared.

As regards plans and programmes which are likely to have significant effects on the environment in another Member State, the Member State in whose territory the plan or programme is being prepared must consult the other Member State(s).

The SEA and EIA differ as follows:

- the SEA requires the environmental authorities to be consulted at the screening stage;
- scoping (i.e. the stage of the SEA process that determines the content and extent of the matters to be covered in the SEA report to be submitted to a competent authority) is obligatory under the SEA;
- the SEA requires an assessment of reasonable alternatives (under the EIA the developer chooses the alternatives to be studied);
- under the SEA Member States must monitor the significant environmental effects of the implementation of plans/programmes to identify unforeseen adverse effects and undertake appropriate remedial action.
- the SEA obliges Member States to ensure that environmental reports are of a sufficient quality.

The SEA Directive applies to a wide range of public plans and programmes. An SEA is mandatory for plans/programmes which are:

- prepared for agriculture, forestry, fisheries, energy, industry, transport, waste/ water management, telecommunications, tourism, town & country planning or land use and which set the framework for future development consent of projects listed in the EIA Directive.

OR

- have been determined to require an assessment under the Habitats Directive.

Broadly speaking, for the plans/programmes not included above, the Member States have to carry out a screening procedure to determine whether the plans/programmes are likely to have significant environmental effects. If there are significant effects, an SEA is needed. The screening procedure is based on criteria set out in Annex II of the Directive.

Annex 2: UK and Norway National Legislation

2.1. United Kingdom

On 1 January 2021, the UK ceased to be a member of the European Union. Whilst much of the European legislation already in place through the national legal system was rolled over, with two new key pieces of national legislation introduced: the Fisheries Act 2020 and the Environment Act 2021.

2.1.1. Fisheries Act 2020

From 1 January 2021, the UK took responsibility for fisheries management within its Exclusive Economic Zone (EEZ) when it left the European Union and the Common Fisheries Policy ceased to apply.

The Fisheries Act 2020 established the legal commitment to fish sustainably, to achieve maximum sustainable yield for each stock and to regulate fishing in order to protect the marine environment. The Fisheries Act notes that the UK will take an ecosystem-based approach to ensure that any negative impacts on marine ecosystems are minimised and, where possible, reversed, and to ensure that incidental catches of sensitive species are minimised and, where possible, eliminated. The Fisheries Act provides the framework for fisheries management in UK waters, including the need for a Joint Fisheries Statement and the development of Fisheries Management Plans.

Fisheries management is devolved to each of the UK administrations. The Joint Fisheries Statement (JFS) outlines the strategies adopted across the nations to meet sustainability and other objectives of the Fisheries Act. The development of the JFS and subsequent Fisheries Management Plans provide an important opportunity for fisheries and marine conservation science communities to work together with neighbouring states to positively shape the future management of fisheries. The JFS reiterates the commitment to minimise and where possible eliminate the bycatch of sensitive species such as cetaceans.

Since March 2022 all fishing vessel licences now contain a mandatory requirement to report the occurrence any marine mammal bycatch within 48 hours of return to port.

2.1.2. Environment Act 2021

Environment Act includes marine and coastal environments within its definition of environment. However, thereafter there is no explicit consideration. The Act introduces the concept of legally binding targets against which implementation progress can be measured. The UK Government have indicated that these targets will include marine biodiversity through a focus on protected areas, resource productivity and plastic pollution. Key indicators for the marine environment, however, are those originally developed under the UK Marine Strategy.

2.1.3. UK Marine Strategy

The UK Marine Strategy and the achievement of Good Environmental Status also emphasises the urgent need to reduce bycatch. The UK Governments have agreed that the same indicators that have been adopted through OSPAR will be utilised in UK waters. This includes indicators M4 Cetacean Abundance and Distribution and M6 marine Mammal Bycatch.

2.1.4. Marine Wildlife Bycatch Mitigation Initiative

The Marine Wildlife Bycatch Mitigation Initiative (BMI) has also been introduced, which outlines how the UK will achieve its ambitions to minimise and, where possible, eliminate the bycatch of sensitive marine species such as harbour porpoise.

The BMI brings together existing work and commits to work that will enable the UK to meet its national and international obligations. Five policy objectives have been identified:

- Improve our understanding of bycatch and entanglement of sensitive marine species through monitoring and scientific research.
- Identify “hotspot” or high-risk areas, gear types and/or fisheries for bycatch and entanglement in the UK in which to focus monitoring and mitigation.
- Develop, adopt and implement effective measures to minimise and, where possible, eliminate bycatch and entanglement of sensitive marine species.
- Identify and adopt effective incentives for fisheries to implement bycatch and entanglement mitigation measures.
- Work with the international community to share best practice and lessons learned to contribute to the understanding, reduction and elimination of bycatch and entanglement globally.

Addressing bycatch whilst simultaneously ensuring productive commercial fisheries is complex and challenging. There is no “one size fits all” approach, instead there needs to be focused, local solutions for each fishery where the bycatch of marine wildlife has been identified as an issue.

This initiative acknowledges the need for fisheries policy authorities to work closely with stakeholder groups across the actions identified to minimise and, where possible, eliminate bycatch of sensitive marine species. These stakeholders include the fishing industry, Non-Government Organisations (NGOs), scientists, experts and innovators.

2.2. Norway

2.2.1 The Nature Diversity Act 2009

The purpose of this Act is to protect biological, geological and landscape diversity and ecological processes through conservation and sustainable use, and in such a way that the environment provides a basis for human activity, culture, health and well-being, now and in the future, including a basis for Sami culture.

The Act implements Norway’s various international commitments, including those of the Bern Convention and the Convention on Biological Diversity. The Act makes provisions for species and habitat protection. It’s implementation in the marine environment is supported by fisheries acts.

2.2.2 The Marine Resources Act 2017

This Act makes provision with respect to the management and conservation of marine living resources. The Act also provides rules relative to marine fishing, with the principal responsibility for administration and control being held by Fisheries Directorate.

2.2.3 Pinger Mandate 2020

This mandate requires all gill net vessels operating in the Vestford to use pingers from 1 January 2021. The mandate has been extended to incorporate over 5000 coastal gill net vessels in an effort to ensure that the USA Marine Mammal Protection Act requirements are being met, enabling the continuation of fisheries exports.

Reference

- Aarefjord, H., A. J. Bjørge, C. C. Kinze, and I. Lindstedt. 1995. Diet of the harbour porpoise (*Phocoena phocoena*) in Scandinavian waters. Rep Int Whal Comm Spec Issue 16:211–222.
- Addink, M. J., and C. Smeenk. 1999. The harbour porpoise *Phocoena phocoena* in Dutch coastal waters: analysis of strandings records for the period 1920-1994. *Lutra* **41**:55-79.
- Advice, I. 2016. Bycatch of small cetaceans and other marine animals – review of national reports under Council Regulation (EC) No. 812/2004 and other information. ICES Special Request Advice.
- ASCOBANS. 2009. ASCOBANS Conservation Plan for Harbour Porpoises (*Phocoena phocoena* L.) in the North Sea.
- Bailey, H., B. Senior, D. Simmons, J. Rusin, G. Picken, and P. M. Thompson. 2010. Assessing underwater noise levels during pile-driving at an offshore windfarm and its potential effects on marine mammals. *Marine Pollution Bulletin* **60**:888-897.
- Benke, H., and U. Siebert. 1996. The current status of harbour porpoises (*Phocoena phocoena*) in German waters. International Whaling Commission, SC/47/SM49, Cambridge.
- Bjørge, A., M. Skern-Mauritzen, and M. C. Rossman. 2013. Estimated bycatch of harbour porpoise (*Phocoena phocoena*) in two coastal gillnet fisheries in Norway, 2006–2008. Mitigation and implications for conservation. *Biological Conservation* **161**:164-173.
- Brandt, M. J., A. Diederichs, K. Betke, and G. Nehls. 2011. Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. *Marine Ecology Progress Series* **421**:205-216.
- Brasseur, S., P. J. H. Reijnders, O. D. Henriksen, J. Carstensen, J. Tougaard, J. Teilmann, M. F. Leopold, C. J. Camphuysen, and J. C. D. Gordon. 2004. Baseline data on the harbour porpoise, *Phocoena phocoena*, in relation to the intended windfarm site NSW, in the Netherlands. Report No. 1043, Alterra, Wageningen.
- Camphuysen, K. 2004. The return of the harbour porpoise (*Phocoena phocoena*) in Dutch coastal waters. *Lutra* **47**:135-144.
- Chehida, Y. B., R. Loughnane, J. Thumloup, K. Kaschner, C. Garilao, P. E. Rosel, and M. C. Fontaine. 2021. No leading-edge effect in North Atlantic harbor porpoises: Evolutionary and conservation implications. *Evol Appl* **14**:1588-1611.
- Dähne, M., A. Gilles, K. Lucke, V. Peschko, S. Adler, K. Krügel, J. Sundermeyer, and U. Siebert. 2013. Effects of pile-driving on harbour porpoises (*Phocoena phocoena*) at the first offshore wind farm in Germany. *Environmental Research Letters* **8**:025002.
- Dähne, M., J. Tougaard, J. Carstensen, A. Rose, and J. Nabe-Nielsen. 2017. Bubble curtains attenuate noise from offshore wind farm construction and reduce temporary habitat loss for harbour porpoises. *Marine Ecology Progress Series* **580**:221-237.
- De Pierrepont, J. F., B. Dubois, S. Desormonts, M. B. Santos, and J. P. Robin. 2005. Stomach contents of English Channel cetaceans stranded on the coast of Normandy. *Journal of the Marine Biological Association of the United Kingdom* **85**:1539-1546.
- Deaville, R. 2016. Annual Report for the period 1st January – 31st December 2015. (Contract number MB0111). Available from: <http://ukstrandings.org/csip-reports/>
- Deaville, R. 2018. UK Cetacean Strandings Investigation Programme final contract report to Defra (MB0111 2011-2017).
- Deaville, R., and P. D. Jepson. 2011. UK Cetacean Strandings Investigation Programme. Final report for the period 1st January 2005 – 31st December 2010. (Covering contract numbers CR0346 and CR0364).
- Donovan, G. P., and A. Bjørge. 1995. Harbour Porpoises in the North Atlantic: edited extract from the Report of the IWC Scientific Committee, Dublin 1995. International whaling commission, Special Issue 16:1-25.
- Evans, P. G. H. 1992. Status of cetaceans in British and Irish waters., UK Mammal Society Cetacean Group, Oxford.
- Evans, P. G. H. 2020. European Whales, Dolphins, and Porpoises. Maine Mammal Conservation in Practice. Academic Press, ISBN 9780128190531.
- Evans, P. G. H., P. Anderwald, and M. E. Baines. 2003. UK Cetacean Status Review. Report to English Nature and Countryside Council for Wales. Sea Watch Foundation, Oxford. 160pp.
- Evans, P. G. H., C. A. Carrington, and J. J. Waggitt. 2021. Risk Mapping of Bycatch of Protected Species

- Evans, P. G. H., C. H. Lockyer, C. S. Smeenk, M. Addink, and A. J. Read. 2008. Harbour porpoise *Phocoena phocoena*. Pages 50-55 in S. Harris and D. W. Yalden, editors. Mammals of The British Isles. Handbook 4th Edition. The Mammal Society.
- FAO. 2018. Report of the Expert Workshop on Means and Methods for Reducing Marine Mammal Mortality in Fishing and Aquaculture Operations. FAO Fisheries and Aquaculture Report No.1231. Rome, Italy. <http://www.fao.org/3/19993EN/i9993en.pdf>.
- Fontaine, M. C., S. J. E. Baird, S. Piry, N. Ray, K. A. Tolley, S. Duke, A. Birkun Jr, M. Ferreira, T. Jauniaux, A. Llavona, B. Ozturk, A. A. Ozturk, V. Ridoux, E. Rogan, M. Sequeira, U. Siebert, G. A. Vikingsson, J.-M. Bouquegneau, and J. R. Michaux. 2007. Rise of oceanographic barriers in continuous populations of a cetacean: the genetic structure of harbour porpoises in Old World waters. *BMC Biology* **5**.
- Fontaine, M. C., K. Roland, I. Calves, F. Austerlitz, F. P. Palstra, K. A. Tolley, S. Ryan, M. Ferreira, T. Jauniaux, A. Llavona, B. Öztürk, A. A. Öztürk, V. Ridoux, E. Rogan, M. Sequeira, U. Siebert, G. A. Vikingsson, A. Borrell, J. R. Michaux, and A. Aguilar. 2014. Postglacial climate changes and rise of three ecotypes of harbour porpoises, *Phocoena phocoena*, in western Palearctic waters. *Molecular Ecology* **23**:3306-3321.
- Fontaine, M. C., O. Thatcher, N. Ray, S. Piry, A. Brownlow, N. J. Davison, P. Jepson, R. Deaville, and S. J. Goodman. 2017. Mixing of porpoise ecotypes in southwestern UK waters revealed by genetic profiling. *Royal Society Open Science* **4**:160992.
- Fontaine, M. C., K. A. Tolley, J. R. Michaux, A. Birkun, M. Ferreira, T. Jauniaux, Á. Llavona, B. Öztürk, A. A. Öztürk, V. Ridoux, E. Rogan, M. Sequeira, J.-M. Bouquegneau, and S. J. E. Baird. 2010. Genetic and historic evidence for climate-driven population fragmentation in a top cetacean predator: the harbour porpoises in European water. *Proceedings of the Royal Society B: Biological Sciences*.
- Galatius, A., C. C. Kinze, and J. Teilmann. 2012. Population structure of harbour porpoises in the Baltic region: evidence of separation based on geometric morphometric comparisons. *Journal of the Marine Biological Association of the United Kingdom* **92**:1669-1676.
- Gilles, A., S. Viquerat, E. Becker, K. Forney, S. Geelhoed, J. Haelters, J. Nabe-Nielsen, M. Scheidat, U. Siebert, S. Sveegaard, F. van Beest, R. van Bemmelen, and G. Aarts. 2016. Seasonal habitat-based density models for a marine top predator, the harbor porpoise, in a dynamic environment. *Ecosphere* **76**:e01367. 01310.01002/ecs01362.01367
- Haelters, J., T. Jauniaux, and J. van Gompel. 2002. Increased number of harbour porpoise strandings in Belgium between 1990 and 2001. Poster presented at the 16th Annual Conference of the European Cetacean Society, Liège, Belgium.
- Haelters, J., F. Kerckhof, E. Toussaint, T. Jauniaux, and S. Degraer. 2012. The diet of harbour porpoises bycaught or washed ashore in Belgium, and relationship with relevant data from the strandings database. ASCOBANS North Sea Harbour Porpoise Conservation Plan, Bonn.
- Hammond, P. S., C. Lacey, A. Gilles, S. Viquerat, P. Börjesson, H. Herr, K. Macleod, V. Ridoux, M. B. Santos, M. Scheidat, J. Teilmann, J. Vingada, and N. Øien. 2017. Estimates of cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys. SCANS-III project report 1, 39pp.
- Hammond, P. S., K. Macleod, P. Berggren, D. L. Borchers, M. L. Burt, A. Cañadas, G. Desportes, G. P. Donovan, A. Gilles, D. Gillespie, J. Gordon, L. Hiby, I. Kuklik, R. Leaper, K. Lehnert, M. F. Leopold, P. Lovell, N. Øien, C. G. M. Paxton, V. Ridoux, E. Rogan, F. Samarra, M. Scheidat, M. Sequeira, U. Siebert, H. Skov, R. Swift, M. L. Tasker, J. Teilmann, O. Van Canneyt, and J. A. Vázquez. 2013. Cetacean abundance and distribution in European Atlantic shelf waters to inform conservation and management. *Biological Conservation* **164**:107-122.
- IAMMWG. 2015. Management Units for cetaceans in UK waters (January 2015). JNCC Report No. 547, JNCC Peterborough.
- ICES. 2021. Workshop on estimation of MOrtality of Marine MAMmals due to Bycatch (WKMOMA). ICES Scientific Reports. 3:106. 95 pp. <https://doi.org/10.17895/ices.pub.9257>.
- ICES. 2022. Working Group on Bycatch of Protected Species (WGBYC). ICES Scientific Reports. 4:91. 265 pp. <https://doi.org/10.17895/ices.pub.21602322>.

-
- ICES Advice. 2014. Bycatch of small cetaceans and other marine animals – review of national reports under Council Regulation (EC) No. 812/2004 and other published documents. ICES Special Request Advice. 8pp.
- ICES Advice. 2021a. Bycatch of protected, endangered, and threatened species (PETS). ICES Advice Ecoregions in the Northeast Atlantic and adjacent seas. Published 2 December 2021. byc.eu – <https://doi.org/10.17895/ices.advice.9335>.
- ICES Advice. 2021b. OSPAR request to estimate bycatch mortality of marine mammals (harbour porpoise *Phocoena phocoena*, common dolphin *Delphinus delphis*, grey seal *Halichoerus grypus*) within the OSPAR maritime area. ICES Special Request Advice Northeast Atlantic ecoregions. Published 2 December 2021. sr.2021.17– <https://doi.org/10.17895/ices.advice.9186>.
- ICES WGBYC. 2020. Report from the Working Group on Bycatch of Protected Species (WGBYC). ICES Scientific Reports. Volume 2, Issue 81. 209 pp.
- ICES WGMME. 2013. Report of the Working Group on Marine Mammal Ecology (WGMME), 4-7 February 2013, Paris, France.
- ICES WGMME. 2014. Report of the Working Group on Marine Mammal Ecology (WGMME), 10-13 March 2014, Woods Hole, Massachusetts, USA. ICES CM 2014/ACOM:27. 234 pp.
- ICES WGMME. 2015. Report of the Working Group on Marine Mammal Ecology (WGMME), 9–12 February 2015, London, UK. ICES CM 2015/ACOM:25. 114pp. ICES CM 2015/ACOM:25. 114 pp.
- IJsseldijk, L. L., K. C. J. Camphuysen, G. O. Keijl, G. Troost, and G. Aarts. 2021a. Predicting Harbor Porpoise Strandings Based on Near-Shore Sightings Indicates Elevated Temporal Mortality Rates. *Frontiers in Marine Science* **8**.
- IJsseldijk, L. L., S. Hensing, A. Mairo, M. T. I. ten Doeschate, J. Treep, J. van den Broek, G. O. Keijl, U. Siebert, H. Heesterbeek, A. Gröne, and M. F. Leopold. 2021b. Nutritional status and prey energy density govern reproductive success in a small cetacean. *Scientific Reports* **11**:19201.
- IJsseldijk, L. L., M. F. Leopold, L. Begeman, M. J. L. Kik, L. Wiersma, M. Morell, E. L. Bravo Rebolledo, T. Jauniaux, H. Heesterbeek, and A. Gröne. 2022. Pathological findings in stranded harbor porpoises (*Phocoena phocoena*) with special focus on anthropogenic causes. *Frontiers in Marine Science* **9**.
- IJsseldijk, L. L., M. T. I. ten Doeschate, A. Brownlow, N. J. Davison, R. Deaville, A. Galatius, A. Gilles, J. Haelters, P. D. Jepson, G. O. Keijl, C. C. Kinze, M. T. Olsen, U. Siebert, C. B. Thøstesen, J. van den Broek, A. Gröne, and H. Heesterbeek. 2020. Spatiotemporal mortality and demographic trends in a small cetacean: Strandings to inform conservation management. *Biological Conservation* **249**:108733.
- IMR-NAMMCO. 2019. Norwegian Institute of Marine Research and the North Atlantic Marine Mammal Commission. Report of Joint IMR/NAMMCO International Workshop on the Status of Harbour Porpoises in the North Atlantic. Tromsø, Norway.
- Jansen, O. E. 2013. Fishing for Food, Feeding ecology of harbour porpoises *Phocoena phocoena* and white-beaked dolphins *Lagenorhynchus albirostris* in Dutch waters. Wageningen University, Wageningen.
- Jauniaux, T., D. Petitjean, C. Brenez, M. Borrens, L. Brosens, J. Haelters, T. Tavernier, and F. Coignoul. 2002. Post-mortem findings and causes of death of harbour porpoises (*Phocoena phocoena*) stranded from 1990 to 2000 along the coastlines of Belgium and Northern France. *J Comp Pathol.* **126**:243-253.
- Jepson, P. D., P. M. Bennett, R. Deaville, C. R. Allchin, J. R. Baker, and R. J. Law. 2005. Relationships between polychlorinated biphenyls and health status in harbour porpoises (*Phocoena phocoena*) stranded in the United Kingdom. *Environmental Toxicology and Chemistry* **24**:238-248.
- Jepson, P. D., R. Deaville, J. L. Barber, À. Aguilar, A. Borrell, S. Murphy, J. Barry, A. Brownlow, J. Barnett, S. Berrow, A. A. Cunningham, N. J. Davison, M. ten Doeschate, R. Esteban, M. Ferreira, A. D. Foote, T. Genov, J. Giménez, J. Loveridge, Á. Llavona, V. Martin, D. L. Maxwell, A. Papachlimitzou, R. Penrose, M. W. Perkins, B. Smith, R. de Stephanis, N. Tregenza, P. Verborgh, A. Fernandez, and R. J. Law. 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Scientific Reports* **6**:18573.
- Jung, J. L., E. Stephan, M. Louis, E. Alfonsi, C. Liret, F. G. Carpentier, and S. Hassani. 2009. Harbour porpoises (*Phocoena phocoena*) in north-western France: aerial survey, opportunistic sightings and strandings monitoring. *Journal of the Marine Biological Association of the UK* **89**:1045-1050.
- Kesselring, T., S. Viquerat, R. Brehm, and U. Siebert. 2017. Coming of age: - Do female harbour

-
- porpoises (*Phocoena phocoena*) from the North Sea and Baltic Sea have sufficient time to reproduce in a human influenced environment? PLoS ONE **12**:e0186951.
- Kiszka, J. J., J. Haelters, and T. Jauniaux. 2004. The harbour porpoise (*Phocoena phocoena*) in the southern North Sea: a come-back in northern French and Belgian waters? Document AC11/Doc.24(P/R) presented at the 11th Advisory Committee Meeting to ASCOBANS, Jastrzebia Góra, Poland, 27 – 29 April, 2004. 4 pp. .
- Lacey, C., A. Gilles, P. Börjesson, H. Herr, K. Macleod, V. Ridoux, M. B. Santos, M. Scheidat, J. Teilmann, S. Sveegaard, J. Vingada, S. Viquerat, N. Øien, and P. S. Hammond. 2022. Modelled density surfaces of cetaceans in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys. Report from SCANS III.
- Lah, L., D. Trense, H. Benke, P. Berggren, P. Gunnlaugsson, C. Lockyer, A. Öztürk, B. Öztürk, I. Pawliczka, A. Roos, U. Siebert, K. Skóra, G. Víkingsson, and R. Tiedemann. 2016. Spatially Explicit Analysis of Genome-Wide SNPs Detects Subtle Population Structure in a Mobile Marine Mammal, the Harbor Porpoise. PLoS ONE **11**:e0162792.
- Lambert, E. 2020. The Feeding Ecology of the Harbour Porpoise *Phocoena Phocoena* L. in a Changing Environment. .
- Laran, S., M. Authier, A. Blanck, G. Doremus, H. Falchetto, P. Monestiez, E. Pettex, E. Stephan, O. Van Canneyt, and V. Ridoux. 2017. Seasonal distribution and abundance of cetaceans within French waters: Part II: The Bay of Biscay and the English Channel. Deep-Sea Research II **14**.
- Law, R. J., J. Barry, J. L. Barber, P. Bersuder, R. Deaville, R. J. Reid, A. Brownlow, R. Penrose, J. Barnett, J. Loveridge, B. Smith, and P. D. Jepson. 2012. Contaminants in cetaceans from UK waters: status as assessed within the Cetacean Strandings Investigation Programme from 1990 to 2008. Marine Pollution Bulletin **64**:1485-1494.
- Leaper, R., and S. Calderan. 2018. Review of methods used to reduce risks of bycatch and entanglements. UNEP/CMS Secretariat, Bonn, Germany. 76pages. CMS Technical Series No. 38.
- Learmonth, J. A., C. D. MacLeod, M. B. Santos, G. J. Pierce, H. Q. P. Crick, and R. A. Robinson. 2006. Potential effects of climate change on marine mammals. Oceanography and Marine Biology: An Annual Review **44**:431-464.
- Leopold, M., and C. J. Camphuysen. 2006. Bruinvisstrandingen in Nederland in 2006: achtergronden, leeftijdsverdeling, sexratio, voedselkeuze en mogelijke oorzaken. IMARES Rep C083/06 and NIOZ Rep 2006-5. Wageningen IMARES and Koninklijk Nederlands Instituut voor Onderzoek der Zee (NIOZ), Texel.
- Leopold, M., F., L. Begeman, J. van Bleijswijk, D. L., L. IJsseldijk, L. , H. Witte, J., and A. Gröne. 2015. Exposing the grey seal as a major predator of harbour porpoises. Proceedings of the Royal Society B: Biological Sciences **282**:20142429.
- Leopold, M., M. Scheidat, M. van den Heuvel-Greve, O. Jansen, A. Beerman, G. Aarts, M. Kottermans, S. Glorius, and S. Bierman. 2011. Aantallen, strandingen en voedsel生态学 van bruinvissen. IMARES rapport BO-11-007-00.
- Leopold, M. F. 2015. Eat or be eaten: porpoise diet studies. Wageningen University, Wageningen, NL.
- Lick, R. R. 1991. Parasites from the digestive tract and food analysis of harbour porpoise (*Phocoena phocoena*) from German waters. In: Evans PGH (ed) European research on cetaceans – 5. European Cetacean Society, Cambridge, p 65–68.
- Lockyer, C. 2003. Harbour porpoises in the North Atlantic: biological parameters. Pages 71-91 in T. Haug, G. Desportes, G. A. Víkingsson, and L. Witting, editors. Harbour porpoises in the North Atlantic. Nammco Scientific Publications Volume 5. The North Atlantic Marine Mammal Commission, Tromso.
- Lusseau, D., L. Kindt-Larsen, and F. M. van Beest. 2023. Emergent interactions in the management of multiple threats to the conservation of harbour porpoises. Sci Total Environ **855**:158936.
- MacLeod, C. D., G. J. Pierce, and M. Begoña Santos. 2007. Starvation and sandeel consumption in harbour porpoises in the Scottish North Sea. Biology Letters **3**:535-536.
- Mahfouz, C., T. Meziane, F. Henry, C. Abi-Ghanem, J. Spitz, T. Jauniaux, T. Bouveroux, G. Khalaf, and R. Amara. 2017. Multi-approach analysis to assess diet of harbour porpoises *Phocoena phocoena* in the southern North Sea. Marine Ecology Progress Series **563**:249-259.
- Martin, A. R. 1996. The diet of harbour porpoises (*Phocoena phocoena*) in British waters. SC/47/SM48. Report of the International Whaling Commission.

-
- Moan, A., and A. Bjørge. 2023. Pingers reduce harbour porpoise bycatch in Norwegian gillnet fisheries, with little impact on day-to-day fishing operations. *Fisheries Research* **259**:106564.
- Mortensen, L. O., and F. Thomsen. 2019. Comparative Study of DEPONS and iPCoD. BSH Cumulative Impact Study 2019. .
- Murphy, S., J. L. Barber, J. A. Learmonth, F. L. Read, R. Deaville, M. W. Perkins, A. Brownlow, N. Davison, R. Penrose, G. J. Pierce, R. J. Law, and P. D. Jepson. 2015. Reproductive failure in UK harbour porpoises *Phocoena phocoena*: Legacy of pollutant exposure? *PLoS ONE* **10**:e0131085.
- Murphy, S., R. J. Law, R. Deaville, J. Barnett, M. W. Perkins, A. Brownlow, R. Penrose, N. J. Davison, J. L. Barber, and P. D. Jepson. 2018. Chapter 1 - Organochlorine Contaminants and Reproductive Implication in Cetaceans: A Case Study of the Common Dolphin. Pages 3-38 in M. C. Fossi and C. Panti, editors. *Marine Mammal Ecotoxicology*. Academic Press.
- Murphy, S., M. A. C. Petitguyot, P. D. Jepson, R. Deaville, C. Lockyer, J. Barnett, M. Perkins, R. Penrose, D. NJ, and C. Minto. 2020. Spatio-Temporal Variability of Harbor Porpoise Life History Parameters in the North-East Atlantic. *Frontiers in Marine Science* **7**.
- Murphy, S., E. H. Pinn, and P. D. Jepson. 2013. The short-beaked common dolphin (*Delphinus delphis*) in the North-eastern Atlantic: distribution, ecology, management and conservation status. Pages 193-280 in R. N. Hughes, D. J. Hughes, and I. P. Smith, editors. *Oceanography and Marine Biology: An Annual Review*, Volume 51. CRC Press.
- Murphy, S., J. Tougaard, P. G. H. Evans, F. Caurant, and P. Hammond. 2019. Area Status Report North Sea. Annex 8. Report of Joint IMR/NAMMCO International Workshop on the Status of Harbour Porpoises in the North Atlantic. Tromsø, Norway. North Atlantic Marine Mammal Commission and the Norwegian Institute of Marine Research.
- Nabe-Nielsen, J., R. M. Sibly, J. Tougaard, J. Teilmann, and S. Sveegaard. 2014. Effects of noise and by-catch on a Danish harbour porpoise population. *Ecological Modelling* **272**:242-251.
- Nabe-Nielsen, J., F. M. van Beest, V. Grimm, R. M. Sibly, J. Teilmann, and P. M. Thompson. 2018. Predicting the impacts of anthropogenic disturbances on marine populations. *Conservation Letters* **11**:e12563.
- Northridge, S. 2018. Bycatch. Pages 149-151 in B. Würsig, J. G. M. Thewissen, and K. M. Kovacs, editors. *Encyclopedia of Marine Mammals*. Third Edition. Academic Press.
- OSPAR. 2017. The Intermediate Assessment 2017. Assessment of the marine environment in OSPAR's waters. <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/>.
- Peltier, H., H. J. Baagøe, K. C. J. Camphuysen, R. Czeck, W. Dabin, P. Daniel, R. Deaville, J. Haelters, T. Jauniaux, L. F. Jensen, P. D. Jepson, G. O. Keijl, U. Siebert, O. Van Canneyt, and V. Ridoux. 2013. The stranding anomaly as population indicator: the case of harbour porpoise *Phocoena phocoena* in North-Western Europe. *PLoS ONE* **8**:e62180-e62180.
- Peltier, H., R. Czeck, W. Dabin, P. Daniel, R. Deaville, J. Haelters, L. L. Ijsseldijk, L. F. Jensen, P. D. Jepson, G. Keijl, M. T. Olsen, U. Siebert, O. Van Canneyt, and V. Ridoux. 2018. Small cetacean mortality as derived from stranding schemes: the harbour porpoise case in the northeast Atlantic. Document SC/67b/HIM05 presented to the IWC Scientific Committee.
- Pierce, G. J., A. Brownlow, P. G. H. Evans, L. Ijsseldijk, K. Kamińska, L. Kessler, S. Murphy, E. Pinn, V. Ridoux, M. P. Simmonds, J. Spitz, K. Stockin, and N. Taylor. 2022. Report of the ASCOBANS Resource Depletion Working Group (August 2022). Report to the 27th Meeting of the Advisory Committee, Online 28-30 September 2022. ASCOBANS/AC27/Doc.2.2.
- Pierce, G. J., M. B. Santos, and S. Cerviño. 2007. Assessing sources of variation underlying estimates of cetacean diet composition: a simulation study on analysis of harbour porpoise diet in Scottish (UK) waters. *Journal of the Marine Biological Association of the UK* **87**:213-221.
- Pinn, E. H., K. Macleod, and M. L. Tasker. 2021. Conservation of transnational species: The tensions between legal requirements and best scientific evidence. *Aquatic Conservation: Marine and Freshwater Ecosystems* **31**:3291-3310.
- Rae, B. B. 1965. The food of the common porpoise (*Phocoena phocoena*). *Journal of Zoology* **146**:114–122.
- Rae, B. B. 1973. Additional notes on the food of the Common porpoise (*Phocoena phocoena*). *Journal of Zoology* **169**:127-131.
- Read, A. J., P. Drinker, and S. Northridge. 2006. Bycatch of Marine Mammals in U.S. and Global Fisheries. *Conservation Biology* **20**:163-169.
- Read, F. L. 2021. Cost-benefit analysis for mitigation measures in fisheries with high bycatch. ASCOBANS Technical Series No. 2. .

-
- Reid, J. B., P. G. H. Evans, and S. P. Northridge. 2003. Atlas of cetaceans distribution in north-west European waters. Joint Nature Conservation Committee (JNCC), Peterborough.
- Robbins, J. R., Bouchet, P.J., Miller, D.L., Evans, P.G.H., Waggitt, J.J., Ford, A., and Marley, S.A. 2022. Shipping in the North-east Atlantic: Increasing concerns for marine conservation. *Marine Pollution Bulletin* **179**:113681.
- Roberts, L., S. Collier, S. Law, and A. Gaion. 2019. The impact of marine vessels on the presence and behaviour of harbour porpoise (*Phocoena phocoena*) in the waters off Berry Head, Brixham (South West England). *Ocean & Coastal Management* **179**:104860.
- Ryan, K., S. Mynott, C. Lyons, T. Clare, E. Day, and C. Bell, et al. . 2022. Hauling Up Solutions 2: Exploring new ways to expand the wildlife bycatch reduction toolkit. Final Workshop Report. 27 pp. Available at: www.cleancatchuk.com/HUS2-Report.
- Santos, M. B. 1998. Feeding ecology of harbour porpoises, common and bottlenose dolphins and sperm whales in the northeast Atlantic. Ph.D. University of Aberdeen.
- Santos, M. B., and G. J. Pierce. 2003. The diet of the harbour porpoise (*Phocoena phocoena*) in the Northeast Atlantic. *Oceanography and Marine Biology: an Annual Review* **41**:355–390.
- Santos, M. B., G. J. Pierce, J. A. Learmonth, R. J. Reid, H. M. Ross, I. A. P. Patterson, D. G. Reid, and D. Beare. 2004. Variability in the diet of harbor porpoises (*Phocoena phocoena*) in Scottish waters 1992-2003. *Marine Mammal Science* **20**:1–27.
- Scheidat, M., A. Gilles, K. H. Kock, and U. Siebert. 2004. Harbour porpoise summer abundance and distribution in the German North and Baltic Seas. working paper AC11/Doc. 8(P)Revision 1 presented to ASCOBANS. Jastrzebia Góra, Poland. 10 pp. .
- Scheidat, M., R. Leaper, M. Van Den Heuvel-Greve, and A. Winship. 2013. Setting Maximum Mortality Limits for Harbour Porpoises in Dutch Waters to Achieve Conservation Objectives. *Open Journal of Marine Science* **3**:133-139.
- Schelling, T., L. J. Van der Steeg, and M. F. Leopold. 2014. The diet of harbour porpoises *Phocoena phocoena* in Dutch waters: 2003-2014. IMARES Report C136/14 <https://edepot.wur.nl/317868>.
- Siebert, U., E. Prenger-Berninghoff, and R. Weiss. 2009. Regional differences in bacterial flora in harbour porpoises from the North Atlantic: environmental effects? *Journal of Applied Microbiology* **106**:329-337.
- Siebert, U., J. Stürznickel, T. Schaffeld, R. Oheim, T. Rolvien, E. Prenger-Berninghoff, P. Wohlsein, J. Lakemeyer, S. Rohner, L. Aroha Schick, S. Gross, D. Nachtsheim, C. Ewers, P. Becher, M. Amling, and M. Morell. 2022. Blast injury on harbour porpoises (*Phocoena phocoena*) from the Baltic Sea after explosions of deposits of World War II ammunition. *Environment International* **159**:107014.
- Siebert, U., K. Tolley, G. A. Vikingsson, D. Olafsdottir, K. Lehnert, R. Weiss, and W. Baumgartner. 2006. Pathological findings in harbour porpoises (*Phocoena phocoena*) from Norwegian and Icelandic Waters. *Journal of Comparative Pathology* **134**:134-142.
- Smeenk, C. 1987. The harbour porpoise *Phocoena phocoena* (L., 1758) in the Netherlands: strandings records and decline. *Lutra* **30**:77-90.
- STECF. 2019. Scientific, Technical and Economic Committee for Fisheries (STECF) - Review of the implementation of the EU regulation on the incidental catches of cetaceans (STECF-19-07). doi 10.2760/64091 JRC117515.
- Stone, C. J., K. Hall, S. Mendes, and M. L. Tasker. 2017. The effects of seismic operations in UK waters: Marine Mammal Observer data. *Journal of Cetacean Research and Management* **16**:71-85.
- Stone, C. J., and M. L. Tasker. 2006. The effects of seismic airguns on cetaceans in U.K. waters. *Journal of Cetacean Research and Management* **8**:255–263.
- Sveegaard, S., A. Galatius, R. Dietz, L. Kyhn, J. C. Koblitz, M. Amundin, J. Nabe-Nielsen, M.-H. S. Sinding, L. W. Andersen, and J. Teilmann. 2015. Defining management units for cetaceans by combining genetics, morphology, acoustics and satellite tracking. . *Global Ecology and Conservation* **3**:839-850.
- Todd, V. L. G., W. D. Pearse, N. C. Tregenza, P. A. Lepper, and I. B. Todd. 1999. Diel echolocation activity of harbour porpoises (*Phocoena phocoena*) around North Sea offshore gas installations. *ICES Journal of Marine Science: Journal du Conseil* **66**:734-745.
- Unger, B., H. Herr, H. Benke, M. Böhmert, P. Burkhardt-Holm, M. Dähne, M. Hillmann, K. Wolff-Schmidt, P. Wohlsein, and U. Siebert. 2017. Marine debris in harbour porpoises and seals from German waters. *Mar Environ Res* **130**:77-84.

-
- van Beest, F. M., L. Kindt-Larsen, F. Bastardie, V. Bartolino, and J. Nabe-Nielsen. 2017. Predicting the population-level impact of mitigating harbor porpoise bycatch with pingers and time-area fishing closures. *Ecosphere* **8**:e01785.
- van den Heuvel-Greve, M. J., A. M. van den Brink, M. J. J. Kotterman, C. J. A. F. Kwadijk, S. C. V. Geelhoed, S. Murphy, J. van den Broek, H. Heesterbeek, A. Gröne, and L. L. Ijsseldijk. 2021. Polluted porpoises: Generational transfer of organic contaminants in harbour porpoises from the southern North Sea. *Science of The Total Environment* **796**:148936.
- WGMME, I. 2019. Working Group on Marine Mammal Ecology (WGMME). ICES Scientific Reports. 1:22. 131 pp. <http://doi.org/10.17895/ices.pub.4980>.
- Wiemann, A., L. W. Andersen, P. Berggren, U. Siebert, H. Benke, J. Teilmann, C. Lockyer, I. Pawliczka, K. Skóra, A. Roos, T. Lyrholm, K. B. Paulus, V. Ketmaier, and R. Tiedemann. 2010. Mitochondrial Control Region and microsatellite analyses on harbour porpoise (*Phocoena phocoena*) unravel population differentiation in the Baltic Sea and adjacent waters. *Conservation Genetics* **11**:195-211.
- Williams, R., M. t. Doeschate, D. J. Curnick, A. Brownlow, J. L. Barber, N. J. Davison, R. Deaville, M. Perkins, P. D. Jepson, and S. Jobling. 2020. Levels of Polychlorinated Biphenyls Are Still Associated with Toxic Effects in Harbor Porpoises (*Phocoena phocoena*) Despite Having Fallen below Proposed Toxicity Thresholds. *Environmental Science & Technology* **54**:2277-2286.
- Williams, R. S., A. Brownlow, A. Baillie, J. L. Barber, J. Barnett, N. J. Davison, R. Deaville, M. ten Doeschate, R. Penrose, M. Perkins, R. Williams, P. D. Jepson, O. Lyashevskaya, and S. Murphy. 2023. Evaluation of a marine mammal status and trends contaminants indicator for European waters. *Science of The Total Environment* **866**:161301.
- Wisniewska, Danuta M., M. Johnson, J. Teilmann, L. Rojano-Doñate, J. Shearer, S. Sveegaard, Lee A. Miller, U. Siebert, and Peter T. Madsen. 2016. Ultra-High Foraging Rates of Harbor Porpoises Make Them Vulnerable to Anthropogenic Disturbance. *Current Biology* **26**:1441-1446.
- Wisniewska, D. M., M. Johnson, J. Teilmann, U. Siebert, A. Galatius, R. Dietz, and P. T. Madsen. 2018. High rates of vessel noise disrupt foraging in wild harbour porpoises (*Phocoena phocoena*). *Proceedings of the Royal Society B: Biological Sciences* **285**: <http://dx.doi.org/10.1098/rspb.2017.2314>.
- Wright, A. J., M. Maar, C. Mohn, J. Nabe-Nielsen, U. Siebert, L. F. Jensen, H. J. Baagøe, and J. Teilmann. 2013. Possible causes of a mass stranding of Harbour porpoise in Danish waters in 2005. *PLoS ONE* **8**(2): e55553. doi:10.1371/journal.pone.0055553.